Fisheries and Oceans Canada | Pêches et Océans Canada
Science | Sciences

**CSAS** | **SCCS**

**Canadian Science Advisory Secretariat** | **Secrétariat canadien de consultation scientifique**

**Research Document 2012/174** | **Document de recherche 2012/174**

**National Capital Region** | **Région de la capitale nationale**

# Risk Assessment for Three Dreissenid Mussels (*Dreissena polymorpha, Dreissena rostriformis bugensis*, and *Mytilopsis leucophaeata*) in Canadian Freshwater Ecosystems

# Évaluation des risques posés par trois espèces de moules dreissénidées (*Dreissena polymorpha, Dreissena rostriformis bugensis* et *Mytilopsis leucophaeata*) dans les écosystèmes d'eau douce au Canada

Thomas W. Therriault[1], Andrea M. Weise[2], Scott N. Higgins[3], Yinuo Guo[1*], and Johannie Duhaime[4]

Fisheries & Oceans Canada

[1]Pacific Biological Station
3190 Hammond Bay Road, Nanaimo, BC V9T 6N7

[2]Institut Maurice-Lamontagne
850 route de la Mer, Mont-Joli, QC G5H 3Z48

[3]Freshwater Institute
501 University Drive, Winnipeg, MB R3T 2N6

[4]Great Lakes Laboratory for Fisheries and Aquatic Sciences
867 Lakeshore Road, PO Box 5050, Burlington, Ontario L7R 4A6

* YMCA Youth Intern

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the Secretariat.

This document is available on the Internet at www.dfo-mpo.gc.ca/csas-sccs/

La présente série documente les fondements scientifiques des évaluations des ressources et des écosystèmes aquatiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

Ce document est disponible sur l'Internet à www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1499-3848 (Printed / Imprimé)
ISSN 1919-5044 (Online / En ligne)
© Her Majesty the Queen in Right of Canada, 2013
© Sa Majesté la Reine du Chef du Canada, 2013

Canada

# TABLE OF CONTENTS


**ABSTRACT** ..... IV

**RÉSUMÉ** ..... V

**INTRODUCTION** ..... 1

RISK ASSESSMENT ..... 1

*Rationale* ..... 1

*Scope and Scale* ..... 1

BIOLOGY ..... 2

*Taxonomy* ..... 2

*Species Descriptions* ..... 4

*Habitat Preferences* ..... 5

*Life History* ..... 7

*Population (Genetic) Structure* ..... 10

*Ecological Impacts* ..... 10

*Potential Interactions with Species At Risk* ..... 12

VECTORS ..... 13

*Primary Invasion* ..... 13

*Secondary Invasion/Dispersal (Spread)* ..... 13

DISTRIBUTION ..... 14

*Native Ranges* ..... 14

*Introduced Ranges* ..... 14

**METHODS AND MATERIALS** ..... 15

RISK ASSESSMENT METHODOLOGY ..... 15

*Determining the Probability of Survival (Habitat Suitability)* ..... 15

*Determining the Probability of Arrival* ..... 16

*Determining the Probability of Invasion* ..... 17

*Defining Impacts and Uncertainty* ..... 17

**RISK ASSESSMENT FOR ZEBRA MUSSEL (*DREISSENA POLYMORPHA*)** ..... 17

STEP 1: DETERMINING THE PROBABILITY OF INVASION ..... 17

*Probability of Survival (Habitat Suitability)* ..... 17

*Probability of Arrival* ..... 18

*Probability of Invasion* ..... 19

STEP 2: DETERMINING THE IMPACTS OF INVASION ..... 19

STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION 20

**RISK ASSESSMENT FOR QUAGGA MUSSEL (*DREISSENA ROSTRIFORMIS BUGENSIS*)** ..... **21**

STEP 1: DETERMINING PROBABILITY OF INVASION ..... 21

*Probability of Survival (Habitat Suitability)* ..... *21*

*Probability of Arrival* ..... *21*

*Probability of Invasion* ..... *22*

STEP 2: DETERMINING THE IMPACTS OF INVASION ..... 22

STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION 23

**RISK ASSESSMENT FOR DARK FALSEMUSSEL (*MYTILOPSIS LEUCOPHEATA*)** ..... **23**

STEP 1: DETERMINING THE PROBABILITY OF INVASION ..... 23

STEP 2: DETERMINING THE IMPACTS OF INVASION ..... 23

STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION 23

**SUMMARY** ..... **23**

**CONCLUSIONS** ..... **26**

**ACKNOWLEDGEMENTS** ..... **27**

**REFERENCES** ..... **27**

**TABLES** ..... **36**

**FIGURES** ..... **62**

**APPENDICES** ..... **84**

**Correct citation for this publication:**

Therriault, T.W., Weise, A.M., Higgins S.N., Guo, S. and Duhaime, J. 2013. Risk Assessment for Three Dreissenid Mussels (*Dreissena polymorpha, Dreissena rostriformis bugensis, and Mytilopsis leucophaeata*) in Canadian Freshwater Ecosystems. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/174 v + 88 p.

## ABSTRACT

An ecological risk assessment for three dreissenid mussel species: the Zebra Mussel (*Dreissena polymorpha*); the Quagga Mussel (*D. rostriformis bugensis*); and the Dark Falsemussel (*Mytilopsis leucopheata*) was conducted for freshwater ecosystems in the western Canadian provinces, Ontario, and Quebec. This risk assessment considered probabilities of survival (habitat suitability) and arrival to 108 Canadian sub-drainages and the ecological impacts associated with these species. The ecological risk associated with both Zebra Mussel and Quagga Mussel invasions across the western provinces and watersheds directly adjacent to the Laurentian Great Lakes and St. Lawrence River was considered high. In contrast, the risk was considered low for most of eastern Ontario and Quebec where calcium concentrations were deemed too low to support large (invasive) populations. Due to the high salinity requirements of Dark Falsemussel, the ecological risk associated with this species was considered low for all Canadian freshwater ecosystems. However, the scope of this risk assessment did not consider coastal estuarine habitats where the ecological risk could be substantially higher. The largest ecological impacts associated with Zebra Mussel and Quagga Mussel were negative impacts on biota that inhabit the pelagic (offshore) zones of lakes or rivers (e.g., losses in productivity for phytoplankton, zooplankton, and planktivorous fishes), and to unionid mussels (severe declines in abundance and biodiversity).

## RÉSUMÉ

Une évaluation des risques écologiques posés par trois espèces de moules dreissénidées : la moule zébrée (*Dreissena polymorpha*), la moule quagga (*D. rostriformis bugensis*) et la moule d'Amérique (*Mytilopsis leucopheata*) a été menée pour les écosystèmes d'eau douce des provinces de l'Ouest canadien, de l'Ontario et du Québec. La présente évaluation des risques s'est penchée sur les probabilités de survie (habitats propices) et d'arrivée de ces espèces dans 108 sous-bassins versants canadiens ainsi que leurs impacts écologiques. D'après l'évaluation, les risques écologiques liés aux invasions de la moule zébrée et de la moule quagga dans les provinces et les bassins-versants de l'Ouest situés directement à côté des Grands Lacs laurentiens et du fleuve Saint-Laurent seraient élevés. En revanche, on a jugé que le risque était faible pour la majeure partie de l'Ontario et du Québec, où les concentrations de calcium ont été considérées comme trop faibles pour soutenir de grandes populations (envahissantes). En raison des exigences élevées de la moule d'Amérique en matière de salinité, on considère que le risque écologique lié à cette espèce est faible pour l'ensemble des écosystèmes d'eau douce au Canada. Toutefois, les habitats côtiers estuariennes, où le risque écologique pourrait être beaucoup plus élevé, ne faisaient pas l'objet de la présente évaluation. Les impacts écologiques les plus importants de la moule zébrée et de la moule quagga sont les effets négatifs sur le biote vivant dans les zones pélagiques (extracôtières) des lacs et des rivières (p. ex., pertes de productivité du phytoplancton, du zooplancton et des poissons planctonophages) et sur les unionidés (déclins considérables de l'abondance et de la biodiversité).

# INTRODUCTION

## RISK ASSESSMENT

### Rationale

Non-indigenous species (NIS) pose an enormous risk to native biodiversity and ecosystem function, especially biodiversity (e.g., Sala et al., 2000, Dextrase and Mandrak, 2006). The invasion cycle of arrival, survival, establishment, and spread of NIS will continue in Canada as it has elsewhere around the world. For example, the number of known introduced species continues to increase in the Great Lakes (e.g., Ricciardi et al., 2006) and along the coasts (e.g., Levings et al., 2002). However, not all NIS are equivalent in terms of their ecological impacts and it is expected that a handful of these species have had disproportionally high ecological and socio-economic impacts associated with their invasions. Having the ability to identify these highest risk invaders (ideally prior to arrival) and focus limited resources on these species is critical for resource managers.

The Canadian Council of Fisheries and Aquaculture Ministers Aquatic Invasive Species Task Group (2004) defined invasive alien species as "those harmful alien species whose introduction or spread threatens the environment, the economy or society, including human health". To guide management actions, a risk assessment can be used to identify higher risk invaders, the important vectors for introduction and/or spread, and the potential impacts if introduced. Due to their proximity to freshwater ecosystems in western Canada, and their well described ecological and economic impacts on invaded ecosystems (e.g., Higgins et al., 2008), the western provinces of Canada (British Columbia, Alberta, Saskatchewan, and Manitoba) requested that Fisheries and Oceans Canada (DFO) conduct a formal risk assessment for *Dreissena polymorpha* (Zebra Mussel) and *D. rostriformis bugensis* (Quagga Mussel). Due to subsequent interest, the geographic scope was expanded to include freshwater ecosystems of Ontario and Quebec. During 2009, a closely related species known as the Dark Falsemussel (*Mytilopsis leucopheata*) was identified on a boat being trailered across western Canada, raising concerns that this species also could pose a risk to Canadian freshwater ecosystems, and so the risk of this species to Canadian freshwater systems also was evaluated.

This document summarizes the results of a risk assessment conducted to evaluate the ecological risk posed by these three non-indigenous dreissenid mussels to Canadian freshwaters and contains information for 108 Canadian sub-drainage basins within the Arctic Ocean, Atlantic Ocean, Gulf of Mexico, Hudson Bay, and Pacific Ocean freshwater drainages (Figure 1). The risk assessment is based upon the most current information available on the distribution, habitat suitability, and ecological impacts for the three dreissenid species evaluated.

### Scope and Scale

There is little doubt that NIS have resulted in a wide variety of social and/or economic impacts and, in some cases, socio-economic risk assessments have been conducted (see Binimelis et al., 2007). The risk assessment presented here for the three dreissenid mussels focuses on the potential ecological risks posed by these invaders, and is consistent with the Canadian Science Advisory Secretariat (CSAS) guidelines for provision of scientific advice to clients (managers). In accordance with the CSAS guidelines socio-economic aspects are not considered in this risk assessment.

Predicting the geographic scale and extent of invasions is complex and includes considerable uncertainties. Given sufficient time (and propagule pressure) the likelihood of an invasion to an

point in the future is not useful for managers making decisions on much shorter timeframes. Thus, the temporal scale of this risk assessment is based on the potential arrival of the three dreissenid mussels to a specific sub-drainage basin (see below) within the next five to ten years. The geographic scope of the risk assessment was originally restricted to freshwater ecosystems in the western provinces of Canada (British Columbia, Alberta, Saskatchewan, and Manitoba) but later expanded to freshwaters of Ontario and Quebec based on subsequent interest from these provinces and their willingness to contribute necessary water quality data. Based on the original criteria, the risk assessment considers only freshwater ecosystems and does not consider coastal marine or estuarine habitats. Within this geographic scope, the risk assessment was conducted at the sub-drainage basin (i.e., secondary watershed) level as defined by the Water Survey of Canada (Atlas of Canada 1,000,000 National Frameworks Data, Hydrology – Drainage areas, version 6, as this allowed characterization of risk for a meaningful but manageable number of spatial units. In total, based on the availability of environmental data, the risk assessment evaluates the ecological risk to 108 of the 184 sub-drainage basins (identified by their 3-digit identifier) within the six provinces (Figure 1). Within each sub-drainage habitat suitability (based on calcium concentrations for Zebra Mussel and Quagga Mussel) can vary between individual freshwater ecosystems. While calcium values ($75^{th}$ percentile) were used to assess the habitat suitability for each sub-drainage, the proportion of sub-sub-drainages within each sub-drainage that fell within each habitat suitability category (very low, moderate, high, very high) was recorded as additional information for managers (Appendices A1 and A2).

## BIOLOGY

### Taxonomy

The taxonomic classification of the three dreissenid species considered in this risk assessment is provided below. Information is based on the following literature: Marelli and Gray (1983); Rosenberg and Ludyanskiy (1994); Therriault et al., (2004); Rosenberg and Huber (2011a); and Rosenberg and Huber (2011b).

- Phylum Mollusca
- Class Bivalvia
- Subclass Heterodonta
- Order Veneroida
- Superfamily Dreissenoidea
- Family Dreissenidae
- Subfamily Dreisseninae
    - Genus Congeria
    - Genus Dreissena
        - *Dreissena polymorpha* (Pallas, 1771) *(valid)*
        - Synonyms
            - *Mytilus polymorpha* (Pallas, 1771)
            - *Dreissena andrusovi* (Andrusov, 1897)

- *Dreissena aralensis* (Andrusov, 1897)
- *Dreissena arnouldi* (Locard, 1893)
- *Dreissena bedoti* (Locard, 1893)
- *Dreissena belgrandi* (Locard, 1893)
- *Dreissena complanata* (Locard, 1893)
- *Dreissena curta* (Locard, 1893)
- *Dreissena eximia* (Locard, 1893)
- *Dreissena küsteri* (Dunker, 1855)
- *Dreissena locardi* (Locard, 1893)
- *Dreissena lutetiana* (Locard, 1893)
- *Dreissena magnifica* (Locard, 1893)
- *Dreissena obtusecarinata* (Andrusov, 1897)
- *Dreissena occidentalis* (Locard, 1893)
- *Dreissena paradoxa* (Locard, 1893)
- *Dreissena polymorpha var. lacustrina* (Boettger, 1913)
- *Dreissena recta* (Locard, 1893)
- *Dreissena servaini* (Locard, 1893)
- *Dreissena sulcata* (Locard, 1893)
- *Dreissena tumida* (Locard, 1893)
- *Dreissena ventrosa* (Locard, 1893)
- *Dreissena westerlundi* (Locard, 1893)
- *Mytilus arca* (Kickx, 1834)
- *Mytilus chemnitzii* (Férussac, 1835)
- *Mytilus fluvis* (Gray, 1825)
- *Mytilus hagenii* (Baer, 1826)
- *Mytilus polymorphus fluviatilis* (Pallas, 1771)
- *Mytilus volgensis* (Gray, 1825)
- *Pinna fluviatilis* (Sander, 1780)
- *Tichogonia chemnitzii* (Rossmässler, 1835)

▪ Subspecies:

- *Dreissena polymorpha polymorpha* (Pallas, 1771)
- *Dreissena polymorpha gallandi* (Locard, 1893)
- *Dreissena polymorpha anatolica* (Locard, 1893)

        - *Dreissena rostriformis bugensis* (Andrusov, 1897) (valid)
        - Synonyms:
            - *Dreissena bugensis* (Andrusov, 1897)
            - *Dreissena brardi* (Eichwald, 1885)
            - *Dreissena grimmi* (Andrusov, 1897)
            - *Dreissema distincta* (Andrusov, 1897)
            - *Dreissena rostriformis compressa* (Logvinenko & Starobogatov, 1966)
            - *Dreissena tschaudae var. pontocaspica* (Andrusov, 1897)
            - *Mytilus rostriformis* (Deshayes, 1838)
    - Genus Mytilopsis
        - *Mytilopsis leucophaeata* (Conrad, 1831) (valid)
        - Synonyms:
            - *Congeria leucophaeta* (Conrad, 1831)
            - *Mytilus leucophaeatus* (Conrad, 1831)
            - *Mytilus cochleatus* (Kickx, in Nyst 1835)
            - *Dreissena cochleatus* (Kickx, 1835)
            - *Congeria cochleatus* (Kickx, 1835)
            - *Mytilina cochleata* (Cantraine, 1837)
            - *Dreissena cochleata* (Nyst, 1843)
            - *Tichogonia cochleata* (Dunker, 1853)
            - *Dreissena cumingiana* (Dunker, 1855)
            - *Mytilus americanus* (Recluz, 1858)
            - *Mytilus tenebrosus* (Reeve, 1858)
            - *Dreissena americana* (Fischer, 1858)
            - *Tichogonia americana* (Kuster, 1889)
            - *Congeria cochleata* (Dall, 1898)

It is important to note that there are other *Dreissena* species in addition to those considered here. For example both *Dreissena presbensis* and *Dreissena blanci* have been confused with *Dreissena polymorpha*, especially in the Balkan region where these two species are endemic.

**Species Descriptions**

The Zebra Mussel, *D. polymorpha*, is a mytiliform bivalve about 25-35 mm in size and black with or without white banding (Figures 2 and 3; Ludyanskiy et al., 1993; Effler and Siegfried, 1994; Patterson et al., 2005). As shell patterns are highly variable for this species they should not be used as the definitive feature for identification (Pathy and Mackie, 1993). Viewed from the inside, the shell has a white lining and a large septum (a shelf-like growth close to the beak of the dressenid) to which the

anterior adductor and retractor muscles are attached (Verween et al., 2010). *D. polymorpha* have both inhalant and exhalant siphons that are used for feeding (Benson et al., 2012a). The umbo (or beak) of Zebra Mussel is pointed, the dorsal margins are rounded, contrasting with relatively flattened ventral margins (Pathy and Mackie, 1993; Dermott and Munawar, 1993; Claxton et al., 1998). The flat ventral side distinguishes the Zebra Mussel from other dreissenids.

Although morphologically similar, the Quagga Mussel (*Dreissena rostriformis bugensis*) differs most obviously from *D. polymorpha* in being larger, rounder, and wider (Figures 2 and 3; Lei and Miller, 1994 *in Pelder, 1994*). Further, this species has a convex, rather than flat, ventral surface that causes the Quagga Mussel to topple over if placed on this side. Viewed from the anterior, the Quagga Mussel displays asymmetry in valve shape and can be distinguished by the byssal groove that is located more ventrally and posteriorly than observed in Zebra Mussel. The color (black, cream or white) and band patterns of specimens are highly variable, with some having no bands at all (Marsden et al., 1996). Where bands are visible, they are concentric and tend to fade in color towards the hinge, which is characteristic of Quagga Mussel (Benson et al., 2012b). Additional morphological variability appears to arise due to two morphotypes of Quagga Mussel which are environmentally rather than genetically determined and apparently related to their life in either shallow or deep-water habitats (Peyer et al., 2010). The deep-water morphotypes appear to have "more flattened and dorsal-ventrally compressed" and "more ovular" profile, as well as a more pointed ventral surface (Dermott and Munawar, 1993; Roe and MacIsaac, 1997; Claxton et al., 1998; Peyer et al., 2010).

The Dark Falsemussel or Conrad's Falsemussel, *Mytilopsis leucophaeata,* has a mytiliform shell with byssal threads typical of dreissenids (Figure 4). According to Verween et al. (2010) the shells of the Dark Falsemussel display stripe or "zigzag" patterns as juveniles, much like the Zebra Mussel. Thus, it is extremely difficult (and potentially unreliable) to distinguish the two species at this stage based on shell color/pattern. As adults, the shell of the Dark Falsemussel is marked with concentric rings that may range from cream-like to dark brown making it easier to distinguish from the Zebra Mussel. As juveniles, the shell shape is "elongate and rectangular" but it generally becomes longer and wider in adults, with the ventral surface being more rounded compared to the dorsal side (Verween et al., 2010). In addition, Dark Falsemussel displays valve asymmetry with the right valve larger than the left (Marelli and Gray, 1983). A distinctive feature for this species is the presence of a small tooth (called an apophysis) located near the umbo (or beak) that serves as the origin of anterior retractor muscles. However, this key feature may be absent or underdeveloped in larval stages (Siddall, 1980; Kennedy, 2011). Relative to *Dreissena* spp., *Mytilopsis* spp. have byssal retractor muscles located more posteriorly. Generally, the shell sizes of North American *M. leucophaeata* range from 10-20 mm, depending on the environment but specimens from Florida examined by Siddall (1980) averaged 22 mm.

At the level of the spermatozoa, both Zebra Mussel and Quagga Mussel differ from the Dark Falsemussel in terms of being more tapered and thin at the head portion (Denson and Wang, 1998). Nichols and Black (1994) observed rounder appearance of the D-hinged stage in Quagga Mussel compared to Zebra Mussel. Under laboratory conditions, hybrids between the two species have been spawned but only reared to the D-hinged stage (Nichols and Black, 1994). Although natural hybrids have been suggested they have not been confirmed (but see Voroshilova et al., 2010).

**Habitat Preferences**

*D. polymorpha* primarily inhabit freshwater ecosystems but have been reported from lower salinity, brackish environments as well. Since they have some capacity for osmoregulation they have been found over a wide range of salinities: 0 to 8-12 ppt for adults and 0 to 6 ppt for embryos (Orlova et al., 2005). This species typically is found in lakes, rivers, canals, and estuaries attached to a wide variety

of substrates such as rocks, shellfish, aquatic plants (Pollux et al., 2010 and references therein). When in an aggregate state, mud and sand grains as fine as 0.06-0.5 mm can become bound substrates that serve as suitable settlement substrate for Zebra Mussel, as has been observed in some United States lakes (Beekey et al., 2004). Zebra Mussel generally settle at moderate water depths (4-7 m), but colonization is possible at either shallower or deeper depths (Bially and MacIsaac, 2000; Wacker and Von Elert, 2003). This species tends to be very rare in the profundal zone (>50 m), potentially because this zone is generally characterized by finer sediments and colder (~4°C) temperatures (Dermott and Munawar, 1993; Mills et al., 1993). Kobak (2001) reported Zebra Mussel preferred dimmer habitats, being negatively phototactic. Although Zebra Mussel populations generally are not sustained in low oxygen or hypoxic conditions, some populations have been observed in hypolimnetic and epilimnetic zones with oxygen concentrations of 0.1-11.2 mg/L and 4.2-14.4 mg/L, respectively, while the lower oxygen threshold appears to be 32-40 Torr at 25°C (Benson et al., 2012a). The range of pH tolerance spans 7.4-9.4 with the optimum around 8.5 (Sprung, 1987; Hincks and Mackie, 1997). Although aquatic, dreissenid mussels, like other mussel species, can tolerate some time out of the water. The only information on desiccation tolerance comes from vector transport studies, where Zebra Mussel persisted for 3-10 days on a boat trailer (Ricciardi et al., 1995) and for 13-18 days under higher humidity conditions (McMahon, 2002). As with other bivalves, a significant quantity of calcium is required for shell development and calcium concentrations are considered a major factor in the potential for establishment and development of large populations (Mackie and Claudi, 2010). The calcium thresholds for Zebra Mussel are reported in Table 1. Thresholds for several environmental variables (e.g., temperature, pH, dissolved oxygen, calcium) suggested to limit Zebra Mussel populations is provided in Mackie and Claudi (2010).

Compared to Zebra Mussel, the Quagga Mussel is more energy-efficient and can live and spawn in cooler, more oligotrophic conditions (Roe and MacIsaac, 1997; Baldwin et al., 2002). While widely dispersed in deepwater habitats of the Laurentian Great Lakes, recent studies have indicated Quagga Mussel largely have displaced Zebra Mussel in shallow depths where the latter had previously dominated (Mills et al., 1996; Patterson et al. 2005; Imo et al., 2010). The ability of Quagga Mussel to use a broad range of substrates has been proposed to be a potential fitness advantage over Zebra Mussel in terms of habitat colonization (Peyer et al., 2011). Due to their euryhalinity, Quagga Mussel can be found in both freshwater and brackish water (Orlova et al., 2005) and can occupy both profundal and littoral zones of lakes and rivers (Mills et al., 1996; Baldwin et al., 2002; Stoeckmann, 2003). Quagga Mussel salinity tolerance ranges from 0-6 ppt: 0-4 ppt is supportive of embryonic development and 6 ppt is an upper lethal limit (Rosenberg and Ludyanskiy, 1994; Spidle et al., 1995; Orlova et al., 2005). In Lakes Erie and Ontario, *D. rostriformis bugensis* have been found at depths of up to 60 m and in other Great Lakes up to 130 m (Mills et al., 1993; Mills et al., 1996; Claxton and Mackie, 1998). However, at very shallow depths within the littoral zone of the Great Lakes dreissenids may be exposed to fatal winter conditions (primarily due to high winds, ice scour) especially if attached to mud substrates (Dermott et al., 2003 *in* Orlova et al., 2005). While Zebra Mussel regularly attach to submerged aquatic vegetation, Quagga Mussel generally do not (Diggins et al., 2004); instead preferentially colonizing cobble and gravel (Dermott et al., 2004 in Orlova et al., 2005) or sedimentary surfaces (Mills et al., 1993). The different preferences for attachment on submerged plants (e.g., macrophytes), which can become entangled on recreational boats and boat trailers, may offer one explanation why Zebra Mussel dispersal across the United States has occurred much more rapidly than dispersal of Quagga Mussel (Benson et al., 2012a; Benson et al., 2012b). Further, the tendency for Zebra Mussel to attach to macrophytes offers a clear management tool to limit the dispersal of Zebra Mussel overland. In some states in the United States the transport of macrophytes on boats or trailers is a ticketable offence (e.g., Wisconsin). As with Zebra Mussel, the most widely used environmental criteria to assess the potential for establishment and reproduction of this species is calcium (Table 1). Thresholds for several environmental variables (e.g., temperature, pH, dissolved

oxygen, calcium) suggested to limit Quagga Mussel populations is provided in Mackie and Claudi (2010).

The Dark Falsemussel is the most euryhaline and eurythermal of the three dreissenid mussel species considered here. Typically characterized as an estuarine species, the Dark Falsemussel is especially adapted to living in environments with high sediment loading potentially due to its long incurrent siphon and its ability to close its valves around the byssus (Marelli and Gray, 1983). This species typically is found in oligo- to mesohaline conditions (e.g., 0.5-5 ppt to 6-18 ppt) within their native North American range (Siddall, 1980). However, some studies suggest this species can survive conditions along the entire estuarine gradient (from 0 to 32 ppt) but neither extreme of this gradient appears to support reproduction (Verween et al., 2010). The wide range of salinity tolerances noted in the literature may be correlated with a high efficiency in hyperosmotic regulation (Verween et al., 2010). This presumably would allow Dark Falsemussel to survive in environments that temporarily become unsuitable. This species also appears to have broad temperature tolerances as the climatic ranges reported include temperate, tropical or sub-tropical (Marelli and Gray, 1983). Findings on the species' ability to handle short-term fluctuations in salinity and/or temperature are inconclusive (Wolff, 1969; Kennedy, 2011). Mackie and Claudi (2010) provide thresholds for several environmental variables (e.g., temperature, pH, dissolved oxygen, calcium) suggested to limit Dark Falsemussel populations. There are many suitable substrates for attachment for *M. leucophaeata* including the shells of Eastern Oysters *Crassostrea virginica* (Conrad 1831), between aggregations of the Hooked Mussel *Ischadium recurvum* (Hinkley, 1907), on firm substrates such as pier pilings, sticks, stones, or bottles (Wolff, 1969). In Chesapeake Bay, Dark Falsemussel form successive layers of shell clusters whereby each cohort of the species is being colonized by the next cohort (Kennedy, 2011). The presence of an apophysis in addition to the byssal retractor position in the Dark Falsemussel presumably renders its byssal attachment to substrates superior to that of *Dreissena* spp. when placed in an environment prone to disturbance such as fast water flow or tidal influences (Moore et al., 1991). However, this is inconsistent with attachment experiments performed on the three mussel species (size ~10 mm, distributed in North America), with Dark Falsemussel exhibiting relatively low attachment strength (see references in Table 1 in Kennedy, 2011).

## Life History

### Zebra Mussel (Dreissena polymorpha)

Zebra Mussel are r-strategists with a short maturation time (1-2 years), high fecundity (>1 million eggs per female for each spawning event), a large ability for dispersal aided by a planktonic veliger stage, and the ability for juveniles and adults to attach to a variety of hard surfaces (e.g., boats, trailers, hard shelled animals) that often are transported to different ecosystems (Ludyanskiy et al., 1993). Zebra Mussel are dioecious with external fertilization in the water column. Fertilized eggs develop into veligers within 3-5 days and are free-swimming for up to one month (Pennak, 1989; Mackie and Schloesser, 1996) thereby enhancing natural dispersal capabilities. Maximum larval abundances (unimodal or bimodal) can be measured between April and September (Pollux et al., 2010). Environmental conditions supporting larval development include temperatures between 12-24°C, pH between 7.4-9.4, and calcium concentrations > 12-25 mg/L (Pollux et al., 2010). During the free-living stage, there can be significant long-distance dispersal of mussel veligers downstream (up to 300 km) (Bially and MacIsaac, 2000). Veligers will develop to postveligers before reaching the juvenile stage that is capable of settling, crawling with a foot and attaching to substrates via byssal threads (Pennak, 1989; Benson et al., 2012a). The flat ventral surface of *D. polymorpha* could aid in securing attachment (Claxton et al., 1998) but massive mortality (90-99%) can occur if substrate colonization is unsuccessful/unsuitable, and/or if temperature and oxygen requirements are not met (Stanczykowska, 1977; Mackie and Schloesser, 1996). Further mortalities arise during the veliger stage as dreissenid

veligers are common prey during May–September, the principle growth season (Hecky et al., 2004). Water velocities exceeding 2 m/s may be enough to dislodge mussels (Richman et al., 2011), while other limiting factors include food limitation (Sprung, 1989), and predation by fish larvae, copepods and rotifers (Sprung, 1993). After settling as juveniles, mussels take a few months to reach sexual maturity (Jantz and Neumann, 1998), which coincides with a shell length of approximately 8-10 mm (Benson et al., 2012a). Females typically reproduce during their second year following oogenesis the previous fall, with egg development and release during the spring (Pennak, 1989), which is synchronous with spermatozoa release (Bacchetta et al., 2010). Despite this seasonal cycle, reproduction may continue throughout the year if supported by environmental conditions such as areas of thermal pollution (Pennak, 1989; Mackie and Schloesser, 1996). Several researchers established that 12°C is the lower threshold for spawning (Sprung, 1989; Ram et al., 1996; McMahon, 1996), although Mantecca et al. (2003) reported a spawning population at 10°C. Like other invasive species, *D. polymorpha* is highly fecund and can produce up to $1.6*10^6$ eggs/female/year (Pennak, 1989; Mackie and Schloesser, 1996; Pollux et al., 2010) with mature eggs 30-96 µm in diameter (Pollux et al., 2010). Shell growth occurs at 6–8°C, reaching 1.5–2.0 cm/year during maturation (Benson et al., 2012a). Although actual growth rates are temperature-determined like other bivalves, it appears higher temperatures promote increased growth rates in Zebra Mussel more than in Quagga Mussel (Baldwin et al., 2002).

Zebra Mussel feed like other bivalves using their inhalant siphons and can ingest small particles (0.07–1.0 µm in diameter), but prefer larger ones (Sprung and Rose, 1988). Common prey includes planktonic algae and zooplankton such as tintinnids, rotifers, copepods, and cladocerans (Mackie and Schloesser, 1996; Thorp and Casper, 2003). Bacteria sometimes comprise a significant portion of their diet (Cotner et al., 1995; Silverman et al., 1996). *Dreissena polymorpha* larvae ingest smaller planktonic species (Sprung, 1989) and at times mussel veligers (MacIsaac et al., 1995). Due to their grazing on small zooplankton and phytoplankton, adult Zebra Mussel compete with larger zooplankton, collectively depressing microzooplankton populations and impacting ecosystem structure and function (Wong et al., 2003). One consequence has been a decline in phytoplankton that in turn has allowed diatoms to proliferate (Ackerman et al., 2001). An arguably greater impact stemming from highly efficient filter feeding by dreissenid invaders in North America, a subject of several investigations, has been a shift in aquatic food chain from a predominantly pelagic to benthic one (elaborated in "Impacts" section) (Berg et al., 1996; Zhu et al., 2006). The specific filtration rates of dreissenids may be influenced by several factors, such as: size, concentration and temperature of suspended particles; size and types of algal and bacterial cells; and mussel size (Mackie and Schloesser, 1996; Benson et al., 2012a). In their study of Zebra Mussel and Quagga Mussel food clearance rates, Baldwin et al. (2002) found an increase in clearance rates with doubled food concentrations, but a decrease in the presence of inorganic (clay) particles suggesting they are selective filter feeders. Baldwin et al. (2002) summarized the optimal temperatures for feeding at 10–20°C, and at 24°C, 100mL of water could be filtered hourly by a 20 mm adult mussel (Bunt et al., 1993).

*D. polymorpha* can survive up to 6–9 years (generally 3–4 years) with potential lifespan linked to ambient temperatures where they tend to live shorter lives in warmer lakes (Stanczykowska, 1977; Benson et al., 2012a). Also, multiple cohorts coexist within a population. Dreissenids can reach very high densities (exceeding 1,000,000 individuals $m^{-2}$) in localized areas when conditions are favorable (Ludyanskiy et al., 1993; Effler and Siegfried, 1994; Patterson et al., 2005). This is comparable to populations measured in Holland (summarized by Bij de Vaat, 1991). In North America, aggregate populations may contain around 700,000 individuals $m^{-2}$ (Pollux et al., 2010; Benson et al., 2012a).

Quagga Mussel (Dreissena rostriformis bugensis)

The life history characteristics of the Quagga Mussel are quite similar to those of the Zebra Mussel (see above) and only deviations will be highlighted here. The strategies employed by *D. rostriformis*

*bugensis* for energy-efficiency include lowering its respiration rate at different temperatures such that its metabolic rate is lowered and surplus energy can be invested in physical growth (Stoeckmann, 2003). Although best adapted to utilize phytoplankton as a food source; detritus, bacteria, and a variety of zooplankton species can comprise a portion of their diet (Cotner et al., 1995; Frischer et al., 2000; Roditi et al., 2000; Higgins and Vander Zanden, 2010). As Zebra Mussel and Quagga Mussel have similar diets (Garton et al., 2005), the larger size of Quagga Mussel may confer a competitive advantage over co-occurring Zebra Mussel (Martel et al., 2001 in Garton et al., 2005). However, this advantage may be offset by the more fragile shells of Quagga Mussel that render it more vulnerable to fish predation (Diggins et al., 2004). As noted previously, the Quagga Mussel appears to have a much wider tolerance for cooler temperatures and softer substrates, allowing for the colonization of profundal sediments in the hypolimnia of lakes. At the ecosystem scale, the development of large populations of Quagga Mussel on softer substrates and below the thermocline may allow them to outcompete Zebra Mussel due to a larger pool of free-swimming larvae that can colonize both hard and soft substrates in littoral waters. While at smaller spatial scales Quagga Mussel densities appear similar to Zebra Mussel densities, at the whole ecosystem scale Quagga Mussel densities have the potential to be much larger (due to their ability to colonize soft sediments in both littoral and profundal waters).

Dark Falsemussel (Mytilopsis leucopheata)

The reproductive period of *M. leucophaeata* in North America generally commences in late spring (Menzie, 1980) and in Holland has been observed to span from summer (May/June) to fall (October/November) (Rajagopal et al., 2005; Verween et al., 2005). Optimal spawning conditions include temperatures greater than 12°C (Verween et al., 2005; Kennedy, 2011) and relatively low salinities, but not freshwater (Kennedy, 2011). Egg sizes of the Dark Falsemussel generally are smaller compared to the other dressenid species but as with the other dreissenids fertilization occurs externally, producing larvae within 24 hours (Verween et al., 2010). Similar to other dreissenids, *M. leucophaeata* develops through several stages; first becoming a trochophore (ciliated larva), then a soft-shelled, bilaterally symmetrical veliger (with ciliated velum), a D- or straight-hinged veliger (not ornamented), a veliconcha (ornamented), followed by organogenesis, foot development, and byssal formation that enable the "setting" or benthic stage called pediveliger (Verween et al., 2010). A combination of foot crawling and byssal attachment allows the Dark Falsemussel to find appropriate substrate (Koch, 1989). The final metamorphosis into a juvenile can be accomplished within nine days (see Table 3 in Kennedy, 2011). As larvae, *M. leucophaeata* may be preyed upon by suspension-feeding bivalves, barnacles, jellyfish, and ctenophores; as adults, they are vulnerable to predation by a diversity of estuarine animals like fish, crabs, and waterfowl (Kennedy, 2011 and references therein). Biocide experiments in an effort to control Dark Falsemussel infestations in Europe have found individuals during the breeding season to be more vulnerable to chlorine treatments (Rajagopal et al., 2002).

Dark Falsemussel are filter feeders of phytoplankton (primarily) and zooplankton (Verween et al., 2010). Stomach content analysis of some Florida specimens also has revealed significant portions of inorganic particles (36%) and plant detritus (31%) (Odum and Heald, 1972). In a study by Rajagopal et al. (2005), smaller mussels held at 20°C and 5.6-5.8 ppt exhibited the greatest foot activity. To our knowledge, no study has investigated the filtration rate of *M. leucophaeata* but it is believed to be similar to that of *D. polymorpha* (Verween et al., 2010). Gradual growth occurs throughout the lifetime of this species (Verween et al., 2006 in Verween et al., 2010). Growth rates for Dark Falsemussel have been found to be positively correlated with temperature; negatively correlated with shell size; and not correlated with chlorophyll a concentrations (Verween et al., 2006 in Kennedy, 2011). For example, juveniles found in Amsterdam Harbour averaging 4 mm at the beginning of the summer gradually increased to 24 mm by the fall (Kennedy, 2011 citing (Vorstman, 1933; Schutz, 1969). The average

age of this cohort was only a year and a few months (in Kennedy, 2011), although 2-4 years has been cited as the average lifespan of the Dark Falsemussel (see Verween et al., 2010).

## Population (Genetic) Structure

Zebra Mussel began expanding their range throughout Europe over 200 years ago, substantially earlier than their establishment in the Laurentian Great Lakes of North America (Table 2). This led many to speculate that it was these 'newly' invaded populations in Europe that were the source of individuals transported to the Great Lakes (e.g., Benson et al., 2012a). However, based on genetic similarities, Ricciardi and MacIsaac (2000) suggested the Baltic Sea was a probable origin for Great Lakes Zebra Mussel populations. Several genetic studies have demonstrated similar genetic heterogeneities among invasive and native populations of *D. polymorpha*, suggesting populations were founded and/or maintained from native populations, large size of the founder populations, or frequent genetic mixing as possible mechanisms (Marsden et al., 1995; Brown and Stepien, 2010). Soroka et al. (1997) assessed the genetic structure of a Zebra Mussel population in Poland and found many loci deviating from Hardy-Weinberg equilibrium. It appears that the *D. polymorpha* populations in North America at that time exhibited contrasting results in terms of agreement with Hardy-Weinberg equilibrium (see Soroka et al., 1997). Polish Zebra Mussel populations also seemed to exhibit higher genetic variability than their North American counterparts, presumably due to differential invasion histories, selection pressures, etc. (Soroka et al., 1997). Even though no hybrid populations of Quagga Mussel and Zebra Mussel have been found in the wild, their co-occurrence and overlapping reproductive period in some ecosystems may create conditions that would enhance hybridization potential. Work by Voroshilova et al. (2010) suggests a putative natural hybrid was detected in Rybinsk Reservoir, Russia, using molecular markers.

Ricciardi and MacIsaac (2000) suggested Quagga Mussel populations invasive in North America originated from native populations found in the estuaries of the Southern Bug and Dnieper Rivers from the Black Sea basin. In a microsatellite marker survey of native and invaded ranges of *D. rostriformis bugensis*, Therriault et al. (2005) found no difference in genetic diversity. This finding was consistent with a previous study by Wilson et al. (1999), lending support to the notion that the high genetic diversity of these mussels constituted a factor for their successful invasion histories. Further these researchers also did not find evidence of isolation-by-distance, thereby inferring that jump dispersal may be responsible for secondary transport, especially in North America where transport by recreational boats was inferred. Lastly, Therriault et al. (2005) attribute the lack of genetic differentiation to significant gene flow owing to one or a combination of mechanisms operating in the invasive range: (a) a large inoculum size; (b) rapid population growth; and/or (c) multiple introductions.

Population densities of Dark Falsemussel in newly invaded European habitats generally greatly exceed population densities in their native North American range, a pattern that is perhaps characteristic of newly invaded versus native habitats (Kennedy, 2011). Similarly, Laine et al. (2006) observed up to 28,000 individuals $m^{-2}$ near a power plant cooling water system in Finland. In contrast, the Hudson River has reported density ranges from 1-25 (at 0-3 ppt), 100-200 (5-9 ppt) to 1,000-2,000 (2-6 ppt) individuals $m^{-2}$ (Walton, 1996). There have been no investigations into the genetic diversity of native versus invasive populations of *M. leucophaeata*.

## Ecological Impacts

The impacts of Zebra Mussel and Quagga Mussel on water quality (i.e., environmental impacts) and flora and fauna (i.e., biological impacts) of invaded habitats are well described in the scientific literature. A recent meta-analysis (Higgins and VanderZanden 2010; Higgins *in press*) of the scientific literature and long-term monitoring datasets quantified the mean, variance, and overall structure of

these impacts for lake and river ecosystems across their invaded range (North America and Europe); the general results of this analysis are described below and in Table 3.

It is important to recognize that considerable variation exists in the impacts of Zebra Mussel and Quagga Mussel invasions on water quality and biota of lake and river ecosystems. Despite this variation, consistent patterns in the direction and magnitude of impacts are evident. First, these dreissenid mussels can induce significant and ecologically relevant impacts on water quality and all major trophic levels from sediment bacteria to apex predators (e.g., piscivorous fishes). Rather than being unique, impacts to multiple trophic levels appear a common consequence of dreissenid invasions. Second, evidence from a temporal analysis of dreissenid impacts on several important ecological indictors (e.g., secchi depth, chlorophyll a concentration, total phosphorus concentration) indicated that the magnitude of impacts were pervasive, with no evidence of declining within 20 years post-establishment of these species (Higgins et al. 2011; Higgins *in press*). Third, the direction of impact (i.e., increase, neutral, decrease) at each trophic level largely was dependant on the energy pathway to which the organism belonged. Organisms that were associated with the pelagic-profundal energy pathway (e.g., phytoplankton, zooplankton, profundal zoobenthos, see Figure 5) most often showed declines in biomass or abundance following dreissenid invasions. In contrast, organisms that were associated with the benthic-littoral pathway (e.g., benthic algae, macrophytes, littoral zoobenthos) generally displayed increases in biomass or abundance following dreissenid invasions. Notable exceptions to this general rule were unionid and sphaerrid mussels, which compete for space and/or food with dreissenids. In particular, unionid mussel populations demonstrated large population declines and loss of species following dreissenid invasions (e.g., Gillis and Mackie, 1994; Ricciardi et al., 1997). This response is particularly troublesome since unionid mussels, already imperiled by habitat degradation and over harvesting, are among the most imperiled faunal groups in North America (Ricciardi et al., 1998). Forth, the magnitude of impact on biota within the pelagic-profundal pathway is related to the filtration capacity of the mussel population, which is a function of population density, the size of the ecosystem, and a variety of factors that affect individual filtration rates (e.g., temperature, water velocity, turbidity) and access to the water-column (e.g., depth, vertical and horizontal mixing). Dreissenid densities can vary by several orders of magnitude over space (within and among lakes or rivers) and time (e.g., years), and whole-ecosystem densities largely are unknown. However, impacts appear to scale with ecosystem size with smaller ecosystems showing the largest impacts. For example, mean declines in phytoplankton were highest in rivers (-78% of pre-dreissenid values), followed by small non-stratified lakes (-58% of pre-dreissenid conditions), and deeper stratified lakes (-38% of pre-dreissenid conditions) (Higgins and Vander Zanden, 2010). A similar pattern was found for zooplankton biomass, with mean declines of 76%, 56%, and 40% for rivers, small non-stratified lakes, and stratified lakes, respectively. While the magnitude of impacts tended to increase with decreases in ecosystem size, this does not indicate that large ecosystems are immune from significant impacts. For example, there are widespread reports of significant impacts to water quality and biota within Lakes Erie, Michigan, and Ontario, which are among the largest freshwater ecosystems on the planet. This appears particularly important when key ecosystem components are affected. For example, in Lake Huron, the arrival and spread of Quagga Mussel to deepwater habitats is thought to have caused the collapse of a key diet item (diporeia) for important forage fishes (Lake Whitefish, Alewife), leading to a collapse of these fish species and the multi-million dollar Pacific salmon fishery. Nonetheless, smaller ecosystems such as rivers, shallow non-stratified lakes, and embayments of larger ecosystems (e.g., Bay of Quinte in Lake Ontario) tend to have larger impacts relative to large stratified lakes.

As with lower trophic levels, the impacts of Zebra Mussel and Quagga Mussel invasions on fish health, population status, and community structure appear related to the energetic pathway from which they obtain their food. However, as most fish species are capable of obtaining food from either resource pathway, the largest negative impacts to fish populations likely will occur for species that are obligate planktivores or deepwater benthivores that are unable to efficiently utilize benthic resources in littoral

zones (Mills et al., 2003; Pothoven and Madenjian, 2008; Rennie et al., 2009). The collapse in planktivore and predator communities in Lake Huron (described above) is an example of the response of fish species to dressenids that were unable to efficiently utilize littoral resources after pelagic and profundal resources (e.g., zooplankton, diporeia) declined. In contrast, species that can efficiently utilize benthic-littoral resources would be expected to benefit from Zebra Mussel and Quagga Mussel invasions. In a well documented case study on fish communities in the Hudson River (Strayer et al., 2004), the abundance of pelagic fish species declined by 28%, and the abundance of littoral fish species increased by 97%. There are 14 fish species in North America, and several species of waterfowl, now known to directly use Zebra Mussel or Quagga Mussel as a prey items (Molloy et al., 1997). Some studies have reported that inclusion of dreissenids in fish diets resulted in declines in fish growth or condition (French and Bur, 1996, Hoyle et al., 2008) and it appears that dreissenid shells offer sufficient protection that they often are considered a food source of last resort for most fish species.

As the filter feeding activities of Zebra Mussel and Quagga Mussel remove phytoplankton and other suspended particulate matter from the water column, water clarity often increases substantially following an invasion. Water clarity is a contributing factor to the penetration of solar energy into lakes, affecting the thermocline depth and heat budgets of lakes, and the growth of algae and plants on the lake bottom. Increasing thermocline depths reduces the volume of the hypolimnion, which could increase deepwater anoxia in some lakes and reduce cold water habitat for some fish species. In some systems, such as the lower Laurentian Great Lakes (Lakes Ontario, Erie, and Michigan), dreissenid invasions led to dramatic increases in nuisance blooms of the benthic alga *Cladophora glomerata* (Higgins et al., 2008). These blooms significantly modified benthic habitats; fouled recreational beaches, municipal and industrial water intakes; were associated with increased abundance of indicator bacteria (e.g., *E. coli*) and pathogenic bacteria (e.g., *Salmonella, Shigella, Campylobacter*); were thought to contribute to avian botulism; and were thought to cause localized anoxia to sediments and sediment biota within depositional areas (Higgins et al., 2008). In some locations of the Laurentian Great Lakes (e.g., Saginaw Bay in western Lake Erie) and inland lakes in Michigan, dreissenid invasions led to an increase in toxin producing phytoplankton species, and their toxin (microcystin), even as total phytoplankton biomass declined (Raikow et al., 2004; Knoll et al., 2008). This hepatotoxin is known to affect liver function, and is a concern both for native biota and humans.

Differences in the magnitude of impacts between Zebra Mussel and Quagga Mussel are not well reported in the scientific literature, presumably because Quagga Mussel are more geographically restricted within North America where the majority of studies have been conducted. However, it is reasonable to expect that the magnitude of impacts associated with Quagga Mussel invasions would be higher than that for Zebra Mussel due to their ability to colonize a wider range of habitats and achieve higher population densities at the ecosystem scale. Thus, there is the potential for increased magnitude of impacts of Quagga Mussel invasions, even in habitats already colonized by Zebra Mussel. General descriptions of the direction, magnitude, and probability of Zebra Mussel and Quagga Mussel impacts to various ecosystem parameters are described in the risk assessment methodology section of this document (see below, and Table 3).

**Potential Interactions with Species At Risk**

Schloesser et al. (1998) highlight the impact dreissenid mussels have had on native unionid mussels in the Great Lakes following their introduction, including significant declines in abundance and species diversity at local scales. Where spatial overlap between introduced driessenid mussels and native mollusc species at risk is high then the impact also should be expected to be high. As of November 2011 a number of molluscs have been assigned status by the Committee on the Status of Endangered

Wildlife in Canada (COSEWIC) with 19 being identified as Endangered, three Threatened and six of Special Concern. Of these, several species have the potential to interact with dreissenid mussels (Table 4; COSEWIC, 2011). In Ontario, where Zebra Mussel and/or Quagga Mussel distributions have overlapped distributions of native mussels identified as Species at Risk, the presence of the invasive dreissenid has been identified as a factor limiting recovery. For example, the Eastern Pondmussel (*Ligumia nasuta*) which had a significant distribution overlap with Zebra Mussel saw a reduction of up to 90% following invasion (COSEWIC, 2007). More recently, in British Columbia the threat of Zebra Mussel contributed to the Endangered status assigned to Rocky Mountain Ridged Mussel (*Gonidea angulata*).

Should dreissenid mussels establish high-density populations in freshwater systems beyond their current range in eastern North America, they could potentially affect the fitness of a number of COSEWIC-listed fish species (COSEWIC, 2011; Table 5), depending on the tendency of each species to prey on these mussels or owing to potential changes in productivity associated with trophic changes attributed to invasions in Europe and North America, notably molluscavores and planktivores (see impacts below).

## VECTORS

### Primary Invasion

Of the many potential primary invasion vectors available to aquatic non-indigenous species both *D. polymorpha* and *D. rostriformis bugensis* were introduced to the Laurentian Great Lakes of North America via ballast water (e.g., Hebert et al., 1989; Pathy and Mackie, 1993; Therriault et al., 2004). Arrival as a hull fouling species is less probable for the Zebra Mussel and Quagga Mussel due to relatively long transit times and oceanic environments that exceed salinity tolerances. Pollux et al. (2010) showed that *D. polymorpha* larvae easily can survive ballast water transport for 11-15 days at 12-24°C, which could be extended if optimal conditions were met. In contrast, the salinity tolerant *M. leucopheata* was introduced to the Hudson River and Europe either in ballast water or as a fouling species on ship hulls or other niche areas (Kennedy, 2011).

### Secondary Invasion/Dispersal (Spread)

Many potential vectors of secondary introduction/spread have been identified for dreissenid mussels. Perhaps the most studied is recreational boating (attached to watercraft/trailers or entrained in live-well/bilge/lines) (e.g., Johnson and Padilla, 1996; Orlova et al. 2004; Pollux et al., 2010). Through a program of mandatory boater inspections in the United States as part 100$^{th}$ Meridian Initiative to slow the spread of Zebra Mussel and Quagga Mussel it has become apparent that commercial hauling of recreational boats represent only a small fraction of overland boat transports but represent approximately one-half of fouled boats that were intercepted (www.agri.idaho.gov/Categories/Environment/InvasiveSpeciesCouncil/Inspection_Stations_ALL.php). Through the same program, in 2011 25 driessenid infested boats were identified with five boats of these boats destined for British Columbia (L.-M. Herborg, B.C. Ministry of the Environment, pers. comm.) and by April 2012 11 infested watercraft were detected. Hence, commercial transport of boats appears a particularly important pathway for long-distance dispersal. Also, boat washing stations appear to have successfully prevented dreissenid invasions in some Ontario lakes that were otherwise suitable and in close proximity to invaded lakes (G. Mackie, Univ. of Guelph, pers. comm.).

Dreissenid mussels also can spread via natural dispersal (e.g., drift, attachment to wildlife) or other human-mediated activities (e.g., intra-basin ballast water discharge, canal creation, waterway operations, scientific expeditions) (e.g., Johnson and Carlton, 1996; Stoeckel et al., 1997; Jantz and

Neumann, 1998; Schneider et al., 2003; Orlova et al. 2005; Ricciardi 2006). Natural dispersal is especially important for drainages where there is a large lake or reservoir that can act as a source of propagules for downstream locations (e.g., Therriault et al., 2004). Following the arrival of Zebra Mussel to the Great Lakes this species rapidly reached downstream locations along the Mississippi River hundreds of kilometers away (Figure 6) likely enhanced by natural dispersal and a combination of natural and human-mediated dispersal events then allowed this species to inhabit additional river segments (Benson et al., 2012a).

## DISTRIBUTION

### Native Ranges

Both Zebra Mussel and Quagga Mussel are native to the Ponto-Caspian Region of Eastern Europe. The Zebra Mussel is considered native to the Black Sea basin, including the Sea of Azov (Mills et al., 1996) while the Quagga Mussel is native to the Dnieper and Bug Limans of the Black Sea basin (Van der Velde et al., 2010; Therriault and Orlova, 2010 (referencing Andrusov, 1897; Kharchenko, 1995)). The Dark Falsemussel is native to the Gulf of Mexico and Atlantic coast of the United States (Marelli and Gray, 1983) but is generally rare in its native range (Kennedy, 2011).

### Introduced Ranges

The Zebra Mussel has an extensive freshwater introduced range as a result of an invasion history that dates back to the late 18$^{th}$ century in Russia (see Table 1). Initially spreading north through the Dnieper and Volga River tributaries (Stanczykowska, 1977) this species continues to spread in European waters with only Norway and Iceland escaping Zebra Mussel introductions thus far. This species arrived in the Laurentian Great Lakes of North America in the mid-1980s and has spread extensively around the Great Lakes basin and along the Mississippi River and its tributaries since that time (Figure 6). Although this species has established populations west of the continental divide in the United States, populations have not been reported yet in western Canada (Manitoba, Saskatchewan, Alberta, or British Columbia).

The Quagga Mussel also has invaded parts of Europe and North America. In Europe, this species invaded waterways in the Caspian basin, most notably the Volga River system (Orlova et al., 2004; Therriault et al., 2004) but also Ukraine (Zhulidov et al., 2005), Hungary (van der Velde and Platvoet, 2007), Germany, Netherlands, and Romania (ISSG, 2012). While the Zebra Mussel rapidly expanded through the eastern United States, the Quagga Mussel has largely remained restricted to the Great Lakes basin. More recently, long range overland transport of Quagga Mussel to several western states has occurred (Figure 7). However, similar to Zebra Mussel, Quagga Mussel populations have not been reported from western Canada.

Dark Falsemussel has invaded brackish waters of the North Sea including coastal waters in France, Belgium, Germany, the Netherlands and more recently England, Finland, the Black Sea, and Spain (Table 6). This species has been reported outside its native range in the United States, but not within Canadian waters (Table 6).

# METHODS AND MATERIALS

## RISK ASSESSMENT METHODOLOGY

Figure 8 presents the flow diagram of the risk assessment process used here for dreissenid mussels. The risk to the environment (ecological risk ONLY as socio-economic risk was not assessed here) posed by a NIS is a combination of the probability of invasion and the impacts to the environment due to that invasion within the risk assessment area. The probability of invasion (Step 1, Figure 8) is determined by the probability of arrival and probability of survival of the NIS. The probability of survival represents the overlap between the physiological requirements/tolerances of a potential invader and environmental conditions in the risk assessment area and is determined here by considering calcium requirements combined with potential temperature limitations. Specifically, we employ calcium concentration thresholds linked to the potential for dreissenid mussels to survive, reproduce, and reach population densities that would be considered invasive within each of the 108 sub-drainages where we had sufficient data. In addition to suitable habitat required for survival, potential invaders must have a mechanism to reach the risk assessment area – the potential for arrival. Here we consider the probability of arrival to be a function of propagule pressure, primarily human-mediated activities determined using a Human Footprint Index (described in the following sections), and the proximity to potential source populations of dreissenid mussels. Spread of a NIS following initial establishment is a function of additional suitable habitat and secondary dispersal vectors and pathways within the risk assessment area. Given the spatial scale of this assessment (i.e., sub-drainages) and limited site-specific data, we did not determine explicitly the probability of secondary spread within each sub-drainage. The rapid expansion of these species across North America and Europe indicates that human-mediated activities are highly likely to re-distribute dreissenids within sub-drainages after their initial arrival. Further, by employing the 75$^{th}$ percentile in available calcium concentrations per sub-drainage (see below) this approach suggests within sub-drainages multiple locations of suitable habitats exist for secondary survival. The impacts to the environment are determined in Step 2 and may include, but are not limited to, impacts on biodiversity, trophic disruption, and habitat alteration or destruction. In Step 3, the probability of invasion is combined with the impacts to the environment to obtain the risk to the environment using a heat matrix (Figure 9). Detailed methodology for each of these steps is described in the following sections.

### Determining the Probability of Survival (Habitat Suitability)

We used reported calcium thresholds to characterize the probability of survival (habitat suitability). Although several environmental variables (e.g., temperature, pH, dissolved oxygen, calcium) may limit successful mollusc invasions (e.g., Mackie and Claudi 2010), Whittier et al. (2008) and Neary and Leach (1992) also used calcium concentration as the primary factor determining Zebra Mussel and Quagga Mussel risk. For Zebra Mussel we defined four probability categories ranging from very low to very high (no "low" category) based on species biology while for the less studied Quagga Mussel we defined three categories (no "low" or "moderate" categories) (Table 1). Only when calcium concentrations are very low (< 12 mg/L) do dreissenid mussels fail to establish. Water quality data were provided by the provinces of British Colombia, Alberta, Manitoba, Saskatchewan, Ontario, and Quebec (Table 7). We extracted calcium data for each sampling station or for Ontario we converted alkalinity to calcium following Mackie and Claudi (2010). Data were selected to represent the most recent sampling year. We calculated the 75$^{th}$ percentile calcium value (sensu Whittier et al., 2008) for each of the 108 sub-drainages within the six provinces. Using the 75$^{th}$ percentile ensures the sub-drainage value is determined by the majority of sites within the sub-drainage and is less influenced by a few, divergent locations that might be less representative of the sub-drainage. For sub-drainages with <5 sampling sites, uncertainty is higher and this uncertainty decreases with increased sampling.

In addition to calcium requirements, published literature suggests that water temperature could be limiting for Zebra Mussel populations. Thus, in order to determine the probability of survival we applied a temperature based correction factor to the calcium concentration scores for Zebra Mussel. Following the relationship identified by Strayer et al. (1991) between water temperature and air temperature in the warmest quarter and data from Mackie and Claudi (2010), available air temperature data from Bioclim 10 (http://www.worldclim.org/) were used to lower the probability of survival in watersheds that have suitable calcium concentrations but are considered too cold to support large Zebra Mussel populations. Thus, the correction for temperature employed here was as follows:

Limiting: air temperature in the warmest quarter $<10^{\circ}C$

Probability of survival reduced by 1 category

Potentially or Not Limiting: air temperature in the warmest quarter $\geq 10^{\circ}C$

Probability of survival not changed

Information on the temperature tolerances of Quagga Mussel suggests they are capable of reproduction in cold hypolimnetic waters of the Laurentian Great Lakes. While it is probable that the development of large populations of Quagga Mussel is reduced below some temperature threshold, we did not have sufficient information to include this factor in our analysis and the northern limits of this species in Canada are therefore unknown. For these reasons the probability of survival (habitat suitability) for Quagga Mussel was not corrected for temperature.

The probability of survival (habitat suitability) was based on calcium requirements tempered by temperature requirements (for Zebra Mussel but not Quagga Mussel) and model outputs fell within five probability categories ranging from very low to very high.

**Determining the Probability of Arrival**

The probability of arrival was defined here as a function of propagule pressure and proximity to an invaded habitat (Figure 8). The inclusion of propagule pressure incorporates the understanding that the transport of Zebra Mussel and Quagga Mussel between habitats is associated with human activities (e.g., trailering of recreational boats). We employed the Human Footprint Index (Sanderson et al., 2002; Appendix A3) as a proxy for propagule pressure. This index is a composite factor of human influence corrected by biome type (http://sedac.ciesin.columbia.edu/wildareas/) that integrates data of land use, urbanization, population density, transportation networks and other human activities that are known to facilitate species invasions (Ficetola et al., 2007; Liu et al., 2011). In order to estimate propagule pressure per sub-drainage, mean scores of the Human Footprint Index were binned according to their natural (Jenks) data breaks into five categories ranging from very low to very high (Table 8).

The probability of arrival was considered to be influenced by the proximity to an invaded habitat, similar to studies that utilize a gravity modelling approach (e.g., Leung et al., 2004). We used information on the current distribution of Zebra Mussel and Quagga Mussel invasions across Canada and the United States (Benson et al., 2012a; 2012b) to calculate a proximity correction factor that adjusted propagule pressure scores (Table 8). Watersheds containing an invaded lake or river, or watersheds either directly adjacent to or within two watersheds of those with an invaded site, were considered to have a very high risk of invasion, and propagule pressure scores were increased by 1 category in the calculation of the probability of arrival (Table 8). Since Zebra Mussel dispersal via natural downstream drift can be substantial (see timing of dispersal along the Mississippi River in Figure 6; Benson et al.,

2012a), propagule pressure scores also were increased by 1 category for sub-drainages downstream of known dreissenid infestations along major rivers. With increasing distance away from invaded habitats, the relative risk of arrival should decrease such that propagule pressure scores were not adjusted (Table 8).

### Determining the Probability of Invasion

The probability of invasion was considered to be a function of the probability of survival (habitat suitability) and the probability of arrival (Figure 8). In this analysis the two components were considered to be equally weighted, and thus were averaged to obtain the probability of invasion for each of the 108 sub-drainages unless calcium was below the required threshold. Since a minimum level of calcium must be available to allow dreissenid mussels to develop their shells (survive and reproduce), should this minimum threshold not be attained then the probability of survival and successful population establishment will be very low. Thus, if the probability of survival was scored as "very low", then the probability of invasion also was scored as "very low".

### Defining Impacts and Uncertainty

To ensure consistency when determining expected impacts on specific ecological endpoints we define five categories for each impact, ranging from very high to very low (Table 9). Similarly, to ensure uncertainty is characterized in a standardized way, we provide an explicit definition of each category also ranging from very high to very low, based on the quality of information available (Table 9).

## RISK ASSESSMENT FOR ZEBRA MUSSEL (*DREISSENA POLYMORPHA*)

### STEP 1: DETERMINING THE PROBABILITY OF INVASION

### Probability of Survival (Habitat Suitability)

#### Calcium Suitability

Most sub-drainages in Manitoba, Saskatchewan, Alberta, eastern British Columbia, and the Great Lakes basin have calcium concentrations that could easily support Zebra Mussel populations at high to very high levels (Figure 10). In contrast, most sub-drainages on the Canadian Shield through central and northwestern Ontario and Quebec have very low calcium concentrations as do sub-drainages along the west coast of British Columbia and in parts of northern Saskatchewan (Figure 10). In fact, these calcium concentrations are considered below the threshold required for Zebra Mussel to survive in these sub-drainages. However, localized calcium concentrations (not assessed here) could be more (or less) favorable for survival. Further, at the scale of the risk assessment conducted here there can be considerable intra-sub-drainage variability as evidenced by the range of actual calcium concentration values. To provide a measure of uncertainty in available calcium data, the percentage of data points that fall into each of the calcium tolerance bins is provided (Table 10). This variability is the greatest source of uncertainty when projecting calcium habitat suitability to the sub-drainage spatial scale used in this risk assessment.

#### Temperature Tolerance

A few sub-drainages in northwestern British Columbia and northern Quebec have temperatures that would be considered limiting to Zebra Mussel (Figure 11). Much of British Columbia, Alberta,

Saskatchewan, Manitoba, Ontario, and Quebec have temperatures that are not limiting for Zebra Mussel (Figure 11).

Probability of Survival (Habitat suitability)

Sub-drainages that have the highest probability of survival of Zebra Mussel as determined from calcium concentrations corrected for potential temperature limitations are located in Manitoba, Saskatchewan, Alberta, eastern British Columbia, and the Great Lakes basin (Figure 12, Table 10). Sub-drainages on the Canadian Shield through central and northwestern Ontario and Quebec have very low probability of survival as do sub-drainages along the west coast of British Columbia and in parts of northern Saskatchewan (Figure 12, Table 10). It is important to note that localized conditions (not assessed at this spatial scale) could be more (or less) suitable for Zebra Mussel survival.

**Probability of Arrival**

Propagule Pressure (Human Footprint Index)

Propagule pressure as determined using the Human Footprint Index is very high around the Great Lakes/St Lawrence River (Figure 13). This index is high through southern Manitoba and southern Saskatchewan extending into southern Alberta (Figure 13). Radiating out from these higher indexed areas are bands of moderate to low indexed sub-drainages from British Columbia through to Quebec (Figure 13). The lowest indexed sub-drainages exist across northern Saskatchewan, northern Manitoba, and northwestern Ontario (Figure 13).

Although sub-drainages in British Columbia have a moderate to low level of propagule pressure as determined by the Human Footprint Index, in 2011 five dreissenid fouled boats were intercepted destined for British Columbia. Although it is unknown how many boats are destined for Canadian watersheds at any given time, this ancillary information suggests even sub-drainages classified as moderate to low have the potential to receive propagules and thus are not immune from potential dreissenid mussel introductions.

Proximity to Invaded Habitats

Consistent with Zebra Mussel being first introduced into the Great Lakes and spreading into the Mississippi River and its major tributaries (Figure 6), most infested watersheds are either around the Great Lakes or in the mid-western United States (Figure 14). This creates a ring of sub-drainages in northwestern Ontario, southern Manitoba and Saskatchewan that are in very close proximity to known Zebra Mussel populations. The proximity then diminishes for Canadian sub-drainages moving west across the prairies and into British Columbia.

Zebra Mussels have been confirmed in the Red River system on the United States side of the border (Figure 6). Thus, the probability of arrival for sub-drainages downstream of this location including Lake Winnipeg and the Nelson River are considered very high due to the potential natural dispersal abilities of this species (discussed above).

Zebra Mussel and Quagga Mussel are among the few invaders that are tracked actively with location information maintained by the USGS so reported populations are considered up to date, at least for the United States. However, any jump dispersal event could result in mussels arriving at new locations and this could significantly alter the proximity to invaded habitats for Canadian freshwater sub-drainages. The calculations would need to be updated should this occur.

Probability of Arrival

The probability of Zebra Mussel arrival was high to very high in southern Manitoba and southern Saskatchewan extending into southern Alberta and in south-central Ontario and southern Quebec around the Great Lakes/St. Lawrence watershed (Figure 15, Table 11). The probability of arrival was moderate for central Quebec and Alberta, and southern British Columbia along the border with the United States (Figure 15, Table 11). The probability of Zebra Mussel arrival in the northern part of the Prairie Provinces and northwestern Ontario was low to very low (Figure 15, Table 11).

**Probability of Invasion**

Probability of invasion was calculated by combining the probability of survival and the probability of arrival. Sub-drainages in southern Manitoba (including Lake Winnipeg and the Nelson River), southern Saskatchewan, southern Alberta, and southern Ontario had a very high probability of Zebra Mussel invasion (Figure 16, Table 11). The probability of invasion was moderate to high for most of British Columbia, Alberta, and Ontario not on the Canadian Shield (Figure 16, Table 11). Sub-drainages on the Canadian Shield in northwestern and central Ontario, Quebec, and coastal British Columbia have a very low probability of Zebra Mussel invasion determined primarily by a lack of calcium (Figure 16, Table 11).

## STEP 2: DETERMINING THE IMPACTS OF INVASION

Information on documented and potential impacts on various ecological parameters endpoints was used to determine the final impact score(s) and level of uncertainty (Table 3). This included consideration of published information on impacts of Zebra Mussel and Quagga Mussel invasions throughout Europe and North America (e.g., Higgins and VanderZanden, 2010). Zebra Mussel and Quagga Mussel densities are highly variable in space and time (e.g., can vary by >1 order of magnitude between years and between invaded ecosystems), and in some circumstances local populations can reach extremely high densities – >10,000 individuals per square meter (Ludyanskiy et al., 1993; Effler and Siegfried, 1994; Patterson et al., 2005). Further, evidence from the scientific literature suggests that the magnitude of impacts largely is determined by density, or a combination of density and ecosystem size. Unfortunately, with the exception of calcium, which at low levels may limit dreissenid mussel densities due to their requirement for building shell material, the drivers of mussel density remain poorly understood (e.g., relationship with trophic status) or unknown at a landscape level (e.g., availability of hard substrate). Nonetheless, a recent meta-analysis of Zebra Mussel and Quagga Mussel impacts (Higgins and VanderZanden, 2010) indicates that mean effect sizes for many ecosystem parameters are significant across invaded ecosystems. Thus, while the magnitude of impacts is likely low in ecosystems with minimal calcium concentrations, at concentrations above these lower limits, significant ecosystem impacts are probable. Potential impacts on invertebrate and fish populations, water quality, animal and human health, and biodiversity (including unioid mussels and other species at risk) were considered.

Negative impacts on fish populations were considered moderate to high, with relatively low uncertainty (Table 3). The scientific literature suggests that the direction of the impact (i.e., positive or negative) on fishes is dependant on the resource pathway where fish obtain their prey. For fish species that capitalize on zoobenthos in littoral areas of lakes or rivers, fish abundance will generally increase due to an increased availability of food resources. For fish species that utilize zooplankton or deepwater zoobenthos, and are unable to efficiently switch to littoral zoobenthos, populations generally will decline. The magnitude of impacts appears to be largely dependant on the filtration capacity of the dreissenid mussel population, which is largely a function of dreissenid density and ecosystem size. Also, in general, ecosystem size is a general predictor of the magnitude of impacts with smaller

ecosystems such as rivers having larger impacts than larger ecosystems. However, in some cases where dreissenids negatively impact key dietary items such as *Diporeia sp.* (a deep water amphipod/scud), dramatic impacts on fish populations can occur even in large ecosystems such as the Laurentian Great Lakes. While Zebra Mussel, and to lesser extent Quagga Mussel, have dispersed widely through eastern North America, reports of fish populations crashing are rare. In contrast, reports indicate that littoral zone fishes such as Smallmouth Bass or Yellow Perch have benefited from dreissenid invasions, with increases in population size.

Based on literature accounts of dreissenid invasions elsewhere, impacts on the physio-chemical environment range from low to high with a very low level of uncertainty for most endpoints (Table 3). A large body of literature describes the physical fouling of hard surfaces including water intakes, propellers and ship hulls, docks and piers. This has implications both for the potential ecosystem extent of dreissenid populations but also serves as a reminder about potential secondary dispersal vectors. The physical fouling of industrial intakes and other surfaces has been widely reported in the peer-review literature. Further, in some locations dreissenid invasions have been associated with large blooms of filamentous algae that have clogged water intake screens of nuclear power facilities, requiring costly multi-day shutdowns. Anecdotal reports also have associated recent spates of avian botulism in the Laurentian Great Lakes with dreissenid mussels via toxin bioaccumulation from phytoplankton. Reports of this impact were widespread in the lower Great Lakes, but otherwise geographically restricted and limited temporally (i.e., did not occur in every year).

Since Zebra Mussel impacts were moderate to very high (Table 3) for several ecological endpoints, the impact to the environment was considered to be very high negative in all cases where invasion occurs. This risk category indicates that impacts associated with Zebra Mussel invasion are significant, with a widespread disruption to the factor in question that persists over time or is likely not reversible. The reversibility of Zebra Mussel impacts is not well understood within invaded habitats, however recent literature reports suggest that impacts do not subside within 10-20 years of invasion and potentially much longer (Higgins et al., 2011; Higgins *in press*). The uncertainty of the environmental impacts was considered to be very low (Table 3) given extensive peer reviewed information in the scientific literature (Table 9) on the impacts of this species.

## STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION

To determine the potential risk to the environment, the probability of invasion was crossed with the impacts to the environment associated with an invasion using the heat matrix (Figure 9). For most of western Canada and Ontario, the risk posed by Zebra Mussel was determined to be high (Figure 17, Table 11). In contrast, very low calcium suitability in sub-drainages in northwestern and central Ontario, Quebec, and coastal British Columbia resulted in a low risk posed by Zebra Mussel in these sub-drainages (Figure 17, Table 11).

# RISK ASSESSMENT FOR QUAGGA MUSSEL (*DREISSENA ROSTRIFORMIS BUGENSIS*)

## STEP 1: DETERMINING PROBABILITY OF INVASION

### Probability of Survival (Habitat Suitability)

#### Calcium Suitability

Most sub-drainages in Manitoba, Saskatchewan, Alberta, eastern British Columbia, northern Ontario, and the Great Lakes basin have calcium concentrations that could easily support Quagga Mussel populations at high to very high levels (Figure 18). In contrast, most sub-drainages on the Canadian Shield through central and northwestern Ontario and Quebec have very low calcium concentrations as do sub-drainages along the west coast of British Columbia and in parts of northern Saskatchewan (Figure 18). In fact, these are considered below the threshold required for Quagga Mussel to survive in these sub-drainages but recall that localized conditions (not assessed here) could be more (or less) favorable. At the scale of the risk assessment conducted here there can be considerable intra-sub-drainage variability. To provide a measure of uncertainty in available calcium data, the percentage of data points that fall into each of the calcium tolerance bins is provided (Table 12). This variability is the greatest source of uncertainty when projecting calcium habitat suitability to the sub-drainage spatial scale used in this risk assessment.

Since Quagga Mussel are often found in deeper, colder waterbodies and thus a temperature correction was deemed unnecessary to determine the probability of survival (habitat suitability). Thus, the calcium concentrations represent the probability of survival in this risk assessment for Quagga Mussel.

### Probability of Arrival

#### Propagule Pressure (Human Footprint Index)

Propagule pressure as determined using the Human Footprint Index is very high around the Great Lakes/St Lawrence River (Figure 13). This index is high through southern Manitoba and southern Saskatchewan extending into southern Alberta (Figure 13). Radiating out from these higher indexed areas are bands of sub-drainages with moderate to low Human Footprint Index values from British Columbia through to Quebec (Figure 13). The sub-drainages with the lowest Human Footprint Index values exist across northern Saskatchewan, northern Manitoba, and northwestern Ontario (Figure 13).

Although sub-drainages in British Columbia have a moderate to low level of propagule pressure as determined by the Human Footprint Index, in 2011 five dreissenid fouled boats were intercepted destined for British Columbia. Although it is unknown how many boats are destined for Canadian watersheds at any given time, this ancillary information suggests even sub-drainages classified as moderate to low have the potential to receive propagules and are potentially at risk.

#### Proximity to Invaded Habitats

Consistent with Quagga Mussel also being first introduced into the Great Lakes and spreading into adjacent watersheds (Figure 7) most infested watersheds are either around the Great Lakes or due to several long distance jump-dispersal events, located in the southwestern United States (Figure 19). This creates one ring of sub-drainages in northwestern Ontario and southern Manitoba that are in very close proximity to known Quagga Mussel populations and a second ring emanating from the

southwestern United States extending towards Canadian sub-drainages. Thus proximity is greatest along the Canada-United States border and diminishes for Canadian sub-drainages moving north.

Zebra Mussel and Quagga Mussel are among the few invaders that actively are tracked with location information maintained by the USGS so reported populations are considered up to date, at least for the United States. However, any jump dispersal event could result in mussels arriving at new locations and this could significantly alter the proximity to invaded habitats for Canadian freshwater sub-drainages. Should this occur the calculations would need to be updated.

Probability of Arrival

The probability of Quagga Mussel arrival was very high in southern Manitoba and the Great Lakes into southern Quebec (Figure 20). The probability of arrival of Quagga Mussel to the southern prairies and central Ontario and Quebec was high while the probability of arrival was moderate to low for most of the remaining sub-drainages with the exception of more northern locations where this probability was very low (Figure 20).

**Probability of Invasion**

Probability of invasion was calculated by combining the probability of survival and the probability of arrival. For Quagga Mussel, sub-drainages in the Great Lakes basin, southern Manitoba, southern Saskatchewan and extending into southern Alberta had a very high probability of invasion (Figure 21). Sub-drainages in eastern British Columbia, central and northwestern Alberta, central Saskatchewan and Manitoba, and parts of Ontario had a high probability of Quagga Mussel invasion (Figure 21). Northwestern British Columbia, northeastern Alberta, northern Manitoba and Ontario had sub-drainages with a moderate probability of invasion. Sub-drainages on the Canadian Shield in northwestern and central Ontario, Quebec, and coastal British Columbia have a very low probability of Quagga Mussel invasion determined primarily by a lack of calcium (Figure 21, Table 12).

**STEP 2: DETERMINING THE IMPACTS OF INVASION**

The scientific literature on Quagga Mussel indicates that direction (i.e., positive or negative) of ecological impacts is identical to those of the Zebra Mussel. Further, the magnitude of ecological impacts for Quagga Mussel is at least equal to and potentially higher than those of the Zebra Mussel. These higher impacts appear related to increased densities associated with the colonization of softer substrates and deeper depths. Since Zebra Mussel impacts were considered very highly negative whenever present, Quagga Mussel impacts were considered at the same level – very highly negative (Table 3) in all cases where population establishment occurs. This risk category indicates that impacts associated with Quagga Mussel invasion are significant, with a widespread disruption to the factor in question that persists over time or is likely not reversible. The reversibility of Quagga Mussel impacts is not well understood within invaded habitats, however, recent literature suggests that impacts do not subside within 10-20 years of invasion and potentially much longer (Higgins et al., 2011; Higgins *in press*). The uncertainty of the ecological impacts was considered to be very low (Table 3). This level of uncertainty indicates that there is extensive peer reviewed information in the scientific literature (Table 9) on the impacts of this species.

### STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION

To determine the potential risk, the probability of invasion was crossed with the impacts to the environment using the heat matrix (Figure 9). For most of western Canada and Ontario the ecological risk posed by Quagga Mussel was determined to be high (Figure 22, Table 13). In contrast, very low calcium suitability in sub-drainages in northwestern and central Ontario, Quebec, and coastal British Columbia resulted in a low ecological risk posed by Quagga Mussel (Figure 22, Table 13).

## RISK ASSESSMENT FOR DARK FALSEMUSSEL (*MYTILOPSIS LEUCOPHEATA*)

### STEP 1: DETERMINING THE PROBABILITY OF INVASION

Based on the salinity requirements of Dark Falsemussel (see Habitat Preferences above) the probability of invasion would be considered very low for each of the assessed freshwater sub-drainages. Thus, given the scope of this risk assessment – freshwater Canadian ecosystems – even if calcium and temperature requirements were met, conditions for reproduction and hence establishment, would not be met due to low salinity in most freshwater ecosystems. Verween et al. (2010) and Kennedy (2011) both suggest reproduction is not possible for *M. leucopheata* in freshwater. Thus, the probability of arrival was not explicitly calculated for each of the sub-drainages assessed. Similarly, potential spread of *M. leucopheata* was not determined due to a lack of suitable habitats within sub-drainages that could support establishment of Dark Falsemussel populations. It is important to note that this species likely could encounter suitable habitats in Canadian estuarine systems but this was beyond the scope of the risk assessment presented here.

### STEP 2: DETERMINING THE IMPACTS OF INVASION

For Dark Falsemussel, as for other mollusc invaders, the potential impacts of an invasion are a function of population size. Again, based on salinity tolerances, it is unlikely that this species would reach invasion densities in any Canadian freshwater ecosystem. Hence, the impacts of a Dark Falsemussel invasion are low.

### STEP 3: RISK TO ENVIRONMENT: COMBINING THE PROBABILITY OF INVASION WITH THE IMPACTS OF INVASION

The ecological risk posed by a Dark Falsemussel invasion in any of the Canadian freshwater ecosystems considered here was determined using the heat matrix (Figure 9). Thus, the ecological risk posed by Dark Falsemussel on Canadian freshwater ecosystems is low. However, the risk posed to Canadian estuarine or marine systems could be substantially higher and an additional risk assessment for these waters would be required to determine the actual risk posed.

## SUMMARY

The ecological risk posed to Canadian freshwater ecosystems by Zebra Mussel and Quagga Mussel was high for most watersheds assessed (Figures 17 and 22). Ecological risk was high for most watersheds across the western Canadian provinces and watersheds of Southern Ontario and Quebec

that are adjacent to the lower Great Lakes (Lakes Huron, Erie, and Ontario) or the St. Lawrence River. The remaining watersheds in Ontario and Quebec were generally considered at low risk from Zebra Mussel or Quagga Mussel due to unsuitable calcium concentrations. Similarly, watersheds along the southern coast of British Columbia, including Vancouver Island, were considered at low risk from both species again due to unsuitable calcium concentrations. The ecological risk posed by the Dark Falsemussel to freshwater drainages in Canada was considered low due to the high salinity requirements for reproduction of this species. However, it is important to note that the scope of this risk assessment did not include coastal estuarine ecosystems that have higher salinity levels and are potentially suitable for this species. A separate risk assessment on the susceptibility of estuarine habitats to the Dark Falsemussel, and associated ecological endpoints, would be required to evaluate the risk to these Canadian ecosystems.

The potential ecological impacts associated with Zebra Mussel and Quagga Mussel invasions, and their uncertainty, was assessed for a large number of ecological endpoints; including physical and chemical attributes of freshwater ecosystems, and biota within all major trophic levels. The level of ecological risk, and uncertainty, posed by Zebra Mussel or Quagga Mussel to these endpoints was not homogenous. At a broad level, the invasion of Zebra Mussel and Quagga Mussel is associated with a broad restructuring of energy flow through freshwater ecosystems, with often dramatic declines in the abundance (or biomass) of species associated with the pelagic energy pathway (e.g., phytoplankton, zooplankton, planktivorous fishes; Table 3), and a general increase in the abundance or biomass of organisms associated with benthic-littoral energy pathways (e.g., benthic algae, zoobenthos and fishes in shallow littoral areas). While increases in energy flow through benthic pathways may appear advantageous in that it may offset losses to energy flow through the pelagic energy pathway and provide food resources to fish capable of utilizing energy from either pathway, in some cases the increased energy flow though the benthic pathway has large negative consequences to numerous ecological endpoints. For example, in the lower Laurentian Great Lakes Zebra Mussel and Quagga Mussel increased benthic algal growth to severe nuisance levels, which subsequently increased bacterial counts (of indicator and pathogenic bacteria), and anoxia in nearshore waters. Second, we specifically note a significant risk to Canadian unionid mussels, several of which have been identified as Endangered by COSEWIC (Table 4). As a general rule, the establishment of dreissenids was associated with a 90% decline in unionid mussel abundance within 10 years, with concomitant losses of mussel diversity (e.g., COSEWIC, 2007). Further, the recovery plans for several of these species have identified the continued threat of Zebra/Quagga Mussel as a contributing factor limiting recovery.

Due to the large spatial scales associated with this risk assessment, we used Canadian sub-drainages (i.e., secondary watersheds), rather than individual lakes or rivers. Variability in habitat suitability within sub-drainages was expected, such that the suitability of each watershed to Zebra Mussel or Quagga Mussel invasion was determined using the 75$^{th}$ percentile of calcium concentration data within the sub-drainage. This approach was deemed acceptable since calcium concentrations within lakes and rivers are generally determined by surface geology that is generally consistent at the watershed scale. However, in some cases there was high variability in calcium concentrations between individual lakes or rivers within sub-drainages. Site-specific information for each watershed (e.g., Tables 10 and 12) is provided and could be used in conjunction with local knowledge to better understand the potential risk to individual lakes or rivers within a sub-drainage, particularly where calcium concentrations are variable but suitable, to inform potential management decisions.

While calcium concentrations in lakes and rivers generally are considered to be a prime determinant of habitat suitability for Zebra Mussel and Quagga Mussel; other environmental variables such as alkalinity, carbonate, chlorophyll a, conductivity/TDS, dissolved oxygen, nitrogen, pH, phosphorous, salinity, secchi depth, temperature, total hardness, and turbidity, also have been useful for improving predictions (see Mackie and Claudi, 2010). While such information likely has been collected for

freshwater systems across Canada by various agencies, it is not currently available and accessible at the large spatial scales used in this assessment. An accessible national water quality database, with geo-referenced data, would be useful for future risk assessments for dreissenids and other NIS. Site-specific or region-specific data for these additional variables would improve the accuracy of regional or local risk assessments and should be included where possible (see Mackie and Claudi, 2010). Also, given the documented propensity of dreissenid mussels to rapidly disperse naturally to downstream locations via their veliger stage and understanding of connectivity within sub-drainages would aid understanding potential invasion dynamics. Should large lakes or reservoirs become infested; the continued rain of propagules to downstream locations will pose an ongoing invasion risk.

In addition to data limitations related to environmental variables there are similar data limitations with respect to distributional information on NIS in Canada. Although Zebra Mussel and Quagga Mussel would be considered a high profile species around the Great Lakes where it has had an invasion presence for about 25 years and has been the focus of repeated outreach and education efforts in Canada and the United States, these mussels would have a lower profile in areas not yet invaded and could be overlooked upon initial arrival to novel locations. There is some monitoring and reporting for NIS in Canada. For example, in Ontario where a broad scale plan to survey freshwater lakes on a rotational basis for a variety of NIS exist (J. Brinsmead, ON Ministry of Natural Resources, pers. comm.) and Manitoba where targeted sampling occurs for priority NIS, vectors, and high risk habitats (J. Shead, Manitoba Conservation, pers. comm.). However, in general, NIS monitoring and reporting across Canada lacks standardization. A national approach including standardized monitoring and reporting would provide greater accessibility to NIS information for both researchers and managers and substantially increase our understanding of actual NIS distributions. This information would be invaluable for future risk assessments as it would provide not only positive findings for infested locations that could be updated on a routine basis it also would provide information where NIS have failed to reach (potentially providing clues to why habitats expected to be invaded are not, such as unsuitable calcium concentrations as noted above). In addition to monitoring for NIS, these programs could contribute to environmental data collection needs identified above.

The most important vector for the potential introduction of Zebra Mussel and/or Quagga Mussel to western Canada is the overland transport of recreational and commercial boats originating from invaded habitats in the United States or Ontario. For example, a commercial vessel originating from Lake Mead, Nevada destined for Saskatchewan (April 2012) and fouled with Quagga Mussels recently was intercepted (L. Dalton, Utah AIS Coordinator, pers. comm.). Surveillance programs in the United States identified the vessel and aquatic invasive species coordinators (or representatives) from each state and from Saskatchewan were alerted to the potential threat. The vessel was then quarantined and professionally decontaminated in Utah. Such rapid response powers, access to decontamination units, and inter-jurisdictional cooperation were highly useful in reducing, or in this case mitigating, the risk to Canadian freshwaters highlighting the need for early detection and rapid response plans (see Locke et al., 2011). Information on such incidents and on boater movements in general, would decrease the uncertainty associated with the 'arrival' aspects identified in this risk assessment and could be used in future risk assessments. Also, this information could be used to focus limited resources to high risk vectors and/or locations. Unfortunately this situation is not unique as infested boats have been intercepted over multiple years in British Columbia (L.-M. Herborg, B.C. Ministry of the Environment, pers. comm.) and Manitoba (J. Shead, Manitoba Conservation, pers. comm.) and likely happens much more frequently than is observed. Thus, there is a need to increase rapid response capabilities; including access to decontamination units as such capabilities would assist in reducing the risk of initial invasion. These efforts could be supplemented with 'slow the spread' campaigns if these species invaded western Canada to raise awareness among stakeholders and the public to help mitigate potential spread.

Commercial shipping was identified as the primary invasion vector responsible for delivering Zebra Mussel and Quagga Mussel to the Great Lakes (Hebert et al., 1989) and for delivering Dark Falsemussel to Europe (Kennedy, 2011). Although the scope of this risk assessment was on Canadian freshwater ecosystems with an emphasis on western Canada, the role of this vector should be assessed separately. Given the salinity tolerances of Zebra Mussel and Quagga Mussel they could be introduced to coastal locations outside the Great Lakes by this vector. Further, should Zebra Mussel or Quagga Mussel be introduced to coastal systems, especially along the west coast of North America, they easily could be transported to Canadian waters either in ballast tanks (similar to arrival to Great Lakes) or attached to the hull (ability to close valves in undesirable conditions). As noted with respect to proximity to invaded locations, a coastal introduction would provide a direct link to several sub-drainages in British Columbia, including those associated with the Port of Vancouver and the Fraser River. Further, Dark Falsemussel may have a greater propensity to utilize the commercial shipping vector over Zebra Mussel or Quagga Mussel since this vector was used to invade Europe (Kennedy, 2011).

## CONCLUSIONS

1. The ecological risk posed by Zebra Mussel and Quagga Mussel was high for most of western Canada and sub-drainages around the Great Lakes/St Lawrence River; and low across most of eastern Ontario and Quebec due to low calcium concentrations on the Canadian Shield.

2. The ecological risk posed by Dark Falsemussel to all freshwater sub-drainages considered was low due to higher salinity requirements for reproduction by this species. A risk assessment for *Mytilopsis leucopheata* that includes coastal waters is required to fully address the potential risk posed by this species to Canadian ecosystems. As a brackish water species, the Dark Falsemussel is most likely to arrive and find suitable conditions in Canadian estuarine waters that were not assessed here.

3. The potential ecological impacts of a Zebra Mussel or Quagga Mussel invasion were evaluated for numerous endpoints (socio-economic indicators not considered here). At highest risk were species associated with pelagic zones of lakes or rivers (e.g., expected declines in productivity of phytoplankton, zooplankton, and planktivorous fishes), and for unionid mussels (severe losses to abundance and biodiversity expected).

4. The habitat suitability for Zebra Mussel and Quagga Mussel was determined at the sub-drainage (i.e., secondary watershed) scale based on calcium concentration ($75^{th}$ percentile). It is important to recognize that calcium concentrations of individual lakes or rivers within each watershed could make these specific systems more (or less) suitable than the $75^{th}$ percentile, especially for sub-drainages where calcium concentrations are highly variable or where uncertainty was high due to a lower number of data points (e.g., sub-drainages with < 5 samples).

5. Natural dispersal by dreissenid mussels within connected waterways can not be ignored. If these mussels invade large lakes or reservoirs within sub-drainages, the potential risk to all connected downstream locations increases substantially. Further, should the distribution of Zebra Mussel or Quagga Mussel change, the risk assessment will need to be updated to reflect the change in proximity to invaded locations.

6. A Human Footprint Index was used as a proxy for propagule pressure. Since the overland transport of recreational boats has been identified as a critical arrival/dispersal vector for

dreissenid mussels, a better understanding of this vector specifically would lower uncertainty and provide information on the human-mediated connectedness of sub-drainages within Canada. Also, education and outreach (e.g., postings at boat launches, cleaning stations) and appropriate rapid response capabilities would further reduce the risk associated with this important vector.

7. A national database of geo-referenced water quality data for Canadian aquatic ecosystems (marine, estuarine, and freshwater) is much needed. Although there is published literature on the tolerances of dreissenid mussels to a variety of water quality parameters (e.g., temperature, turbidity, total phosphorus, chlorophyll a), there were insufficient data available and accessible for these variables at the spatial scales needed for modeling (i.e., at a national scale). Such a database would prove invaluable for determining the potential distribution of aquatic invasive species in Canada under current conditions and under future climate scenarios.

## ACKNOWLEDGEMENTS

The authors wish to thank provincial contacts for providing much needed environmental data for British Columbia, Alberta, Saskatchewan, Manitoba, Ontario and Quebec. Dr. Leif-Matthias Herborg, Serge Proulx, and Gilles Fortin provided helpful guidance on ArcGIS and modeling methods. The workshop participants were an essential component of ensuring high quality scientific advice on these potential high risk invaders. DFO's Aquatic Invasive Species program and its Centre of Expertise for Aquatic Risk Assessment provided travel funds.

## REFERENCES

Ackerman, J.D., Loewen, M.R., and Hamblin, P.F. 2001. Benthic-pelagic coupling over a Zebra Mussel reef in western Lake Erie. Limnol. Oceanogr. **46**:892-904.

Bacchetta, R., Mantecca, P. and Vailati, G. 2010. Reproductive behaviour of Zebra Mussels living in shallow and deep water in the South Alps lakes. In: Van der Velde, G., Rajagopal, S., Bij de Vaate, A. (eds) The Zebra Mussel in Europe, Backhuys Publishers, Leiden/Margraf Publishers, Weikersheim. pp. 161-168.

Baldwin B.S., Mayer M.S., Dayton J., Pau N., Mendilla J., Sullivan M., Moore A., Ma A., and Mills E.L. 2002. Comparative growth and feeding in Zebra and Quagga Mussels (*Dreissena polymorpha* and *Dreissena bugensis*): implications for North American lakes. Can. J. Fish. Aquat. Sci. **59**:680-694.

Beekey, M.A., McCabe, D.J., and Marsden, J.E. 2004. Zebra Mussel colonisation of soft sediments facilitates invertebrate communities. Freshw. Biol. **49**:535-545.

Benson, A.J., Raikow, D., Larson, J., and Fusaro, A. 2012a. *Dreissena polymorpha*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL. http://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=5 Revision Date: 2/15/2012 [Accessed January 15, 2012]

Benson, A.J., Richerson, M.M., Maynard, E., Larson, J., and Fusaro, A. 2012b. *Dreissena bugensis*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL. http://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=95 Revision Date: 2/14/2012 [accessed January 15th 2012]

Berg, D.J., Fisher, S.W., and Landrum, P.F. 1996. Clearance and processing of algal particles by Zebra Mussels (*Dreissena polymorpha*). J. Great Lakes Res. **22**:779-788.

Bially, A., and MacIsaac, H.J. 2000. Fouling mussels (*Dreissena* spp.) colonize soft sediments in Lake Erie and facilitate benthic invertebrates. Freshw. Biol. **43**:85-97.

Bij de Vaate, A. 1991. Distribution and aspects of population dynamics of the Zebra Mussel, *Dreissena polymorpha* (Pallas, 1771), in the Lake Usselmeer area (The Netherlands). Oecologia **86**:40-50.

Binimelis, R., Born, W., Monterroso, I., and Rodríguez-Labajos, B. 2007. Socio-economic impact and assessment of biological invasions. In: Nentwig W (ed) Biological Invasions. Ecological Studies 193. Springer-Verlag, Berlin, pp. 331-347.

Brown, J.E., and Stepien, C.A. 2010. Population genetic history of the dreissenid mussel invasions: expansion patterns across North America. Biol. Invasions **12**:3687-3710.

Bunt, C.M., MacIssac, H.J., and Sprules, W.G. 1993. Pumping rates and projected filtering impacts of juvenile Zebra Mussels (*Dreissena polymorpha*) in western Lake Erie. Can. J. Fish Aquat. Sci. **50**:1017-1022.

Canadian Council of Fisheries and Aquaculture Ministers Aquatic Invasive Task Group. 2004. A Canadian Action Plan to Address the Threat of Aquatic Invasive Species. September, 2004. [www.dfo-mpo.gc.ca/Science/enviro/ais-eae/plan/plan-eng.pdf]

Claxton, W.T., and Mackie, G.L. 1998. Seasonal and depth variations in gametogenesis and spawning of *Dreissena polymorpha* and *Dreissena bugensis* in eastern Lake Erie. Can. J. Zool. **76**:2010-2019.

Claxton, W.T., Wilson, A.B., Mackie, G.L., and Boulding, E.G. 1998. A genetic and morphological comparison of shallow- and deep-water populations of introduced dreissenid bivalve *Dreissena bugensis*. Can. J. Zool. **76**:1269-1276.

Cohen, A.N. 2007. Potential Distribution of Zebra Mussels (*Dreissena polymorpha*) and Quagga Mussels (*Dreissena bugensis*) in California Phase 1 Report. A Report for the California Department of Fish and Game, 29 pp.

Cohen, A.N., and Weinstein, A. 2001. Zebra Mussel's Calcium Threshold and Implications for its Potential Distribution in North America. Richmond, CA: San Francisco Estuary Institute. http://www.sfei.org/sites/default/files/2001-Zebramusselcalcium356.pdf [Accessed January 13, 2012].

Conrad, T.A. 1831. Description of fifteen new species of Recent, and three of fossil shells, chiefly from the coast of the United States. J. Acad. Nat. Sci. Philad. **6**:256-268.

COSEWIC. 2007. COSEWIC assessment and status report on the Eastern Pondmussel *Ligumia nasuta* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vi+34 pp.

COSEWIC. 2011. Canadian Wildlife Species at Risk. Committee on the Status of Endangered Wildlife in Canada.

Cotner, J.B, Gardner, W.S., Johnson, J.R., Sada, R.H., Cavaletto, J.F., and Heath, R.T. 1995. Effects of Zebra Mussels (*Dreissena polymorpha*) on bacterioplankton: evidence for both size-selective consumption and growth stimulation. J. Great Lakes Res. **21**:517-528.

Denson, D.R., and Wang, S.Y. 1998. Distinguishing the dark false mussel, *Mytilopsis leucophaeata* (Conrad, 1831), from the non-indigenous Zebra and Quagga Mussels, *Dreissena* spp., using spermatozoan external morphology. Veliger **41**:205-207.

Dermott, R.M., and Munawar, M. 1993. Invasion of Lake Erie offshore sediments by *Dreissena*, and its ecological implications. Can. J. Fish Aquat. Sci. **50**:2298-2304.

Dermott, R., Bonnell, R., Carou, S., Dow, J. and Jarvis, P. 2003. Spatial distribution and population structure of the mussels *Dreissena polymorpha* and *Dreissena bugensis* in the Bay of Quinte Lake Ontario, 1998 and 2000. Can. Tech. Rep. Fish. Aquat. Sci. **2479**:58 pp.

Dextrase, A., and Mandrak, N.E. 2006. Impacts of invasive alien species on freshwater fauna at risk in Canada. Biol Invasions **8**: 13-24.

Diggins, T.P., Weimer, M., Stewart, K.M., Baier, R.E., Meyer, A.E., Forsberg, R.F., and Goehle, M.A. 2004. Epiphytic refugium: are two species of invading freshwater bivalves partitioning spatial resources? Biol. Invasions **6**:83-88.

Drake, J.M, and, Bossenboek, J.M. 2004. The potential distribution of Zebra Mussels in the United States. BioScience **54**:931-941.

Effler, S.W., and Siegfried., C. 1994. Zebra Mussel (*Dreissena polymorpha*) populations in the Seneca River, New York: impact on oxygen resources. Environ. Sci. Tech. **28**:2216-2221.

Ficetola, G.F., Thuiller, W., and Miaud, C. 2007. Prediction and validation of the potential global distribution of a problematic alien invasive species-the American bullfrog. Diversity Distrib. **13**:476-485.

French, J.R.P., and Bur, M.T. 1996. The effect of Zebra Mussel consumption on growth of freshwater drum in Lake Erie. J. Freshwat. Ecol. **11**:283-289.

Frischer, M.E., Nierzwicki-Bauer, S.A., Parsons, R.H., Vathanodorn, K., and Waitkus, K.R. 2000. Interactions between Zebra Mussels (*Dreissena polymorpha*) and microbial communities. Can. J. Fish. Aquat. Sci. **57**:591-599.

Garton, D.W., Payne, C.D., and Montoya, J.P. 2005. Flexible diet and trophic position of dreissenid mussels as inferred from stable isotopes of carbon and nitrogen. Can. J. Fish. Aquat. Sci. **62**:1119-1129.

Gillis, P.L., and Mackie, G.L. 1994. Impact of the Zebra Mussel, *Dreissna polymorpha*, on populations of Unionidae (Bivalvia) in Lake St. Clair. Can. J. Zoo. **72**:1260-1271.

Hebert, P.D., Muncaster, B.W., and Mackie, G.L. 1989. Ecological and genetic studies on *Dreissena polymorpha* (Pallas): a new mollusc in the Great Lakes. Can. J. Fish. Aquat. Sci. **46**:1587-1591.

Hecky, R.E., Smith, R.E.H., Barton, D.R., Guildford, S.J., Taylor, W.D., Charlton, M.N., and Howell, T. 2004. The nearshore phosphorus shunt: a consequence of ecosystem engineering by dreissenids in the Laurentian Great Lakes. Can. J. Fish. Aquat. Sci. **61**:1285-1293.

Higgins, S.N., and VanderZander, M.J. 2010. What a difference a species makes: a meta analysis of dreissenid mussel impacts on freshwater ecosystems. Ecol. Monogr. **80**:179-196.

Hincks, S.S., and Mackie, G.L. 1997. Effects of pH, calcium, alkalinity, hardness, and chlorophyll on the survival, growth, and reproductive success of Zebra Mussel (*Dreissena polymorpha*) in Ontario lakes. Can. J. Fish Aquat. Sci. **54**:2049-2057.

Hinkley, A.A. 1907. Shells collected in northeastern Mexico. Nautilus **21**:68-72.

Hoyle, J.A., Bowlby, J.N., and Morrison, B.J. 2008. Lake whitefish and walleye population responses to dreissenid mussel invasion in eastern Lake Ontario. Aquat. Ecosyst. Health Manage. **11**:403-411.

Imo, M., Seitz, A., and Johannesen, J. 2010. Distribution and invasion genetics of the Quagga Mussel (*Dreissena rostriformis bugensis*) in German rivers. Aquat. Ecol. **44**:731-740.

ISSG. 2012a. *Dreissena polymorpha* (mollusk): Distribution. Global Invasive Species Database. Website: http://www.issg.org/database/species/distribution.asp?si=50&fr=1&sts=sss&lang=EN [Accessed January 16, 2012]

ISSG. 2012b. *Dreissena bugensis* (mollusk): Distribution. Global Invasive Species Database. Website: http://www.issg.org/database/species/distribution.asp?si=918&fr=1&sts=&lang=EN [Accessed January 16, 2012]

ISSG. 2012c. *Mytilopsis Leucophaeata* (mollusk): Distribution. Global Invasive Species Database. Website: http://www.issg.org/database/species/distribution.asp?si=707&fr=1&sts=&lang=EN [Accessed January 16, 2012]

Jantz, D. and Neumann, D. 1998. Growth and reproductive cycle of the Zebra Mussel in the River Rhine as studied in a river bypass. Oecologia **114**:213-225.

Johnson, L.E. and Carlton, J.T. 1996. Post-establishment spread in largescale invasions: Dispersal mechanisms of the Zebra Mussel *Dreissena polymorpha*. Ecology **77**:1686-1690.

Johnson, L.E. and Padilla, D.K. 1996. Geographic spread of exotic species: ecological lessons and opportunities from the invasion of the Zebra Mussel *Dreissena polymorpha*. Biol. Conserv. **78**:23-33.

Kennedy, V.S. 2011a. The invasive Dark Falsemussel *Mytilopsis leucophaeata* (Bivalvia: Dreissenidae): a literature review. Aquat. Ecol. **45**:163-183.

Kennedy, V.S. 2011b. Biology of the uncommon dreissenid bivalve *Mytilopsis leucophaeata* (Conrad, 1831) in central Chesapeake Bay. J. Mollus. Stud. **77**:154-164.

Knoll, L.B., Sarnelle, O., Hamilton, S.K., Kissman, C.E.H., Wilson, A.E., Rose, J.B., and Morgan, M.R. 2008. Invasive zebra mussels (*Dreissna polymorpha*) increase cyanobacterial toxin concentrations in low-nutrient lakes. Can. J. Fish. Aquat. Sci. **65**: 448-455.

Kobak, J. 2001. Light, gravity and conspecifics as cues to site selection and attachment behaviour of juvenile and adult *Dreissena polymorpha* Pallas, 1771. J. Molluscan Stud. **67**:183-189.

Koch, L.M. 1989. *Mytilopsis leucophaeta* (Conrad, 1831) from the upper Mississippi River (Bivalvia: Dreissenidae). Malacol. Data Net. **2**:153-154.

Laine, A.O., Mattila, J., and Lehikoinen, A. 2006. First record of the brackish water dreissenid bivalve *Mytilopsis leucophaeta* in the northern Baltic Sea. Aquat. Invasions **1**:38-41.

Leung, B., Drake, J.M., and Lodge, D.M. 2004. Predicting invasions: propagule pressure and the gravity of allee effects. Ecology **85**:1651-1660.

Levings, C., Kieser, D. Jamieson, G.S., and Dudas, S. 2002. Marine and estuarine alien species in the Strait of Georgia, British Columbia. Alien invaders in Canada's waters, wetlands, and forests. Natural Resources Canada, pp. 111-131.

Liu, X., Guo, Z., Wang, S., and Li, Y. 2011. Increasing potential risk of a global aquatic invader in Europe in contrast to other continents under future climate change. PLoS ONE **6**:1-11.

Locke, A., Mandrak, N.E., and Therriault, T.W. 2011. A Canadian Rapid Response Framework for Aquatic Invasive Species. DFO Can. Sci. Advis. Sec. Res. Doc. 2010/114. vi + 30 p.

Ludyanskiy, M.L., McDonald, D., and Macneill, D. 1993. Impact of the Zebra Mussel, a bivalve invader - *Dreissena polymorpha* is rapidly colonizing hard surfaces throughout waterways of the United States and Canada. BioScience **43**:533-544.

MacIsaac, H.J. Lonnee, C.J. and Leach, J.H. 1995. Suppression of microzooplankton by Zebra Mussels: importance of mussel size. Freshw. Biol. **34**:379-387.

Mackie, G.L., and Claudi, R. 2010. Monitoring and control of macrofouling molluscs in fresh water systems. 2nd Ed. CRC Press, New York. 508 p.

Mackie, G.L., and Schlosser, D.W. 1996. Comparative biology of Zebra Mussels in Europe and North America: an overview, Am. Zool. **36**:244-258.

Mantecca, P., Vailati, G., Garibaldi, L., and Bacchetta, R. 2003. Depth effects on Zebra Mussels reproduction. Malacologia **45**:109-120.

Marelli, D.C., and Gray, S. 1983. Conchological redescription of *Mytilopsis sallei* and *Mytilopsis leucophaeta* of the brackish Western Atlantic (Bivalvia: Dreissenidae). Veliger **25**:185-193.

Marsden, J.E., Spidle, A.P., and May, B. 1995. Genetic similarity among Zebra Mussel populations within North America and Europe. Can. J. Fish. Aquat. Sci. **52**:836-847.

Marsden, J.E., Spidle, A.P., and May, B. 1996. Review of genetic studies of *Dreissena* spp. Am. Zool. **36**:259-270.

Martel, A.L., Baldwin, B.S., Dermott, R.M., and Lutz, R.A. 2001. Species and epilimnion/hypolimnion-related differences in size at larval settlement and metamorphosis in *Dreissena* (Bivalvia). Limnol. Oceangr. **46**:707-713.

McMahon, R.F. 1996. The physiological ecology of the Zebra Mussel, *Dreissena polymorpha*, in North America and Europe. Am. Zool. **36**:339-363.

McMahon, R.F. 2002. Evolutionary and physiological adaptations of aquatic invasive animals: r-selection versus resistance. Can. J. Fish. Aquat. Sci. **59**:1235 -1244.

Menzie, C.A. 1980. The chironomid (Insecta: Diptera) and other fauna of a *Myriophyllum spicatum* L. plant bed in the lower Hudson River. Estuaries **3**:38-54.

Mills, E., Casselman, J.M., Dermott, R. Fitzsimmons, J.D., Gal, T., Holeck, K.T., Hoyle, J.A., Johannssson, O.E., Lantry, B.F., Makarewicz, J.C., Millard, E.S., Munawar, I.F., Munawar, M., O'Gorman,R., Owens, R.W., Rudstam, L.G., Schaner, T., and Stewart, T.J. 2003. Lake Ontario: food web dynamics in a changing ecosystem (1970-2000). Can. J. Fish. Aquat. Sci. **60**:471-490.

Mills, E.L., Dermott, R.M., Roseman, E.F., Dustin, D., Mellina, E., Conn, D.B., and Spidle, A.P. 1993. Colonization, ecology, and population structure of the "Quagga" mussel, (Bivalvia:Dreissenidae) in the lower Great Lakes. Can. J. Fish. Aquat. Sci. **50**:2305-2314.

Mills, E.L., Rosenberg, G., Spidle, A.P., Ludyanskiy, M. Pligin, Y., and May, B. 1996. A review of the biology and ecology of the Quagga Mussel (*Dreissena bugensis*), a second species of freshwater Dreissenid introduced to North America. Am. Zool. **36**:271-286.

Molloy, D.P., A.Y. Karataev, L.E. Burlakova, D.P. Kurandina, and F. Laruelle. 1997. Natural enemies of Zebra Mussels: predators, parasidtes, and ecological competitors. Rev. Fish. Sci. **5**:27-97.

Moore, S.G., O'Neill, C.R. Jr., and MacNeill, D.B. 1991. Ecological perspectives on functional morphologies of *Dreissena polymorpha* and *Mytilopsis leucophaeta*. *Dreissena polymorpha*. Info. Rev. **2**:2-3.

Nichols, S.J., and Black, M.G. 1994. Identification of larvae: the Zebra Mussel (*Dreissena polymorpha*), Quagga Mussel (*Dreissena rosteriformis bugensis*), and Asian clam (*Corbicula fluminea*). Can. J. Zool. **72**:406-417.

Odum, W.E., and Heald, E.J. 1972. Trophic analyses of an estuarine mangrove community. Bull. Mar. Sci. **22**:671-738.

Orlova, M., S. Golubkov, L. Kalinina, and N. Ignatieva. 2004. *Dreissena polymorpha* (Bivalvia: Dreissenidae) in the Neva Estuary (eastern Gulf of Finland, Baltic Sea): Is it a biofilter or source for pollution? Mar. Pollut. Bull. **49**:196-205.

Orlova, M.I., Therriault, T.W., Antonov, P.I. and Shcherbina, G.K. 2005. Invasion ecology of Quagga Mussels (*Dreissena rostriformis bugensis*): a review of evolutionary and phylogenetic impacts. Aquat. Ecol. **39**:401-418.

Pathy, D.A., and Mackie, G.L. 1993. Comparitive shell morphology of *Dreissena polymosrpha, Mytilopsis leucophaeata,* and the 'Quagga' mussel (Bivalvia:Dressenidae) in North America. Can. J. Zool. **71**:1012-1023.

Patterson, M.W.R., Ciborowski, J.J.H., and Barton, D.R. 2005. The distribution and abundance of *Dreissena* species (Dreissenidae) in Lake Erie, 2002. J. Great Lakes Res. **31**:223-237.

Pelder, D. 1994. Distinguishing the Zebra Mussel, Quagga Mussel, and Conrad's False Mussel Based on Shell Morphology. Technical Note ZMR-1-24. US Army Engineers Waterways Experiment Station.

Peyer, S.M., Hermanson, J.C. and Lee, C.E. 2010. Developmental plasticity of shell morphology in Quagga Mussels from shallow and deep-water habitats in the Great Lakes. J. Exp. Biol. **213**:2602-2609.

Peyer, S.M., Hermanson, J.C. and Lee, C.E. 2011. Effects of shell morphology on mechanics of Zebra and Quagga Mussel locomotion. J. Exp. Biol. **214**:2226-2236.

Pollux, B.J.A., Van der Velde and G., Bij de Vaate, A. 2010. A perspective on global spread of *Dreissena polymorpha:* a review on possibilities and limitations. In: Van der Velde, G., Rajagopal, S., Bij de Vaate, A. (eds). The Zebra Mussel in Europe. Backhuys Publishers, Leiden/Margraf Publishers, Weikersheim, pp: 45 - 58

Pothoven, S.A., and Madenjian, C.P. 2008. Changes in consumption by alewives and lake whitefish after dreissenid mussel invasions in Lakes Michigan and Huron. NA J. Fish. Manag. **28**:308-320.

Raikow, D.F., Sarnelle, O., Wilson, A.E., and Hamilton, S.K. 2004. Dominance of the noxious cyanobacterium *Microcystis aeruginosa* in low-nutrient lakes is associated with exotic zebra mussels. Limnol. Oceanogr. **49**:482-487.

Rajagopal, S., Van der Gaag, M., Van der Velde, G., and Jenner, H.A. 2002. Control of brackish water fouling mussel, *Mytilopsis leucophaeta* (Conrad), with sodium hypochlorite. Arch. Environ. Contam. Toxicol. **43**:296-300.

Rajagopal, S., Van der Gaag, M., Van der Velde, G., and Jenner, H.A. 2005. Upper temperature tolerances of exotic brackishwater mussel, *Mytilopsis leucophaeta* (Conrad): an experimental study. Mar. Envir. Res. **60**:512-530.

Ram, J.L., Fong, P.P., and Garton, D.W. 1996. Physiological aspects of Zebra Mussel reproduction: Maturation, spawning and fertilization. Am. Zool. **36**:326-338.

Rennie, M.D., Sprules, W.G., and Johnson, T.B. 2009. Resource switching in fish following a major food web disruption. Oecologia **159**:789-802.

Ribi, G. 1986. Within-lake dispersal of the prosobranch snails, *Viviparus ater* and *Potamopyrgus jenkinsi*. Oecologia **69**:60-63.

Ricciardi, A. 2006. Patterns of invasion in the Laurentian great lakes in relation to changes in vector activity. Divers. Distrib. **12**:425-433.

Ricciardi, A. and MacIssac, H.J. 2000. Recent mass invasions of the North American Great Lakes by Ponto-Caspian species. TREE **15**:62-65.

Ricciardi, A., Serrouya, R., and Whoriskey, F.G. 1995. Aerial exposure tolerance of Zebra and Quagga Mussels (Bivalvia, Dreissenidae) -- implications for overland dispersal. Can. J. Fish. Aquat. Sci. **52**:470-477.

Ricciardi, A., F.G. Whoriskey, and J.B. Rasmussen. 1997. The role of the Zebra Mussel (*Dreissena polymorpha*) in structuring macroinvertebrate communities on hard substrata. Can. J. Fish. Aquat. Sci. **54**: 2596-2608.

Roditi, H.A., Caraco, N.F., Cole, J.J. and Strayer, D.L. 2000. Uptake of dissolved organic carbon and trace elements by Zebra Mussels. Nature **407**:78-80.

Roe, S.L. and MacIssac, H.J. 1997. Deepwater population structure and reproductive state of Quagga Mussels (*Dreissena bugensis*) in Lake Erie. Can. J. Fish. Aquat. Sci. **54**:2428-2433.

Rosenberg, G., and Huber, M. 2011a. *Dreissena bugensis* Andrusov, 1897. Accessed through: World Register of Marine Species at: http://www.marinespecies.org/aphia.php?p=taxdetails&id=505319 [Accessed March 6, 2012]

Rosenberg, G., and Huber, M. 2011b. *Dreissena polymorpha* (Pallas, 1771). Accessed through: World Register of Marine Species at: http://www.marinespecies.org/aphia.php?p=taxdetails&id=181566 [Accessed March 6, 2012]

Rosenberg, G., and Ludyankskiy, M.L. 1994. A nomenclature review of *Dreissena* (Bivalvia: Dreissenidae) with Identification of the Quagga Mussel as *Dreissena bugensis*. Can. J. Fish Aquat. Sci. **51**:1474-1484.

Sala, O.E. Chapin, F.S., Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R. Huber-Sanwald, E., Huenneke, L.F., Jackson, R.B., Kinzig, A., Leemans, R. Lodge, D.M., Mooney, H.A., Oesterheld, M., Poff, N.L., Sykes, M.T., Walker, B.H., Walker, M., and Wall, D.H. 2000. Biodiversity - Global biodiversity scenarios for the year 2100. Science **287**:1770-1774.

Sanderson, E.W., Jaiteh, M., Levy, M.A., Redford, K.H., Wannebo, A.V., and Woolmer, G. 2002. The Human Footprint and the Last of the Wild. BioScience **52**:891-904.

Schloesser, D.W., Kovalak, W.P., Longton, G.D., Ohnesorg, K.L., and Smithee, R.D. 1998. Impact of Zebra and Quagga Mussels (*Dreissena* spp.) on freshwater unionids (Bivalvia: Unionidae) in the Detriot River of the Great Lakes. Am. Midl. Nat. **140**:299-313.

Schneider, D.W., Stoeckel, J.A., Rehmann, C.R., Blodgett, K.D., Sparks, R.E., and Padilla, D.K. 2003. A developmental bottleneck in dispersing larvae: implications for spatial population dynamics. Ecol. Lett. **6**:352-360.

Siddall, S.E. 1980. Early development of *Mytilopsis leucophaeta* (Bivalvia: Dreissenacea). Veliger **22**:378-379.

Silverman, H., Lynn, J.W., Archberger, E.C., Dietz, T.H. 1996. Gill structure in Zebra Mussels: Bacterial-sized particle filtration. Am. Zool. **36**:364-372.

Spidle, A.P., Mills, E.L., and May, B. 1995. Limits to tolerance of temperature and salinity in the Quagga Mussel (*Dreissena bugensis*) and the Zebra Mussel (*Dreissena polymorpha*). Can. J. Fish. Aquat. Sci. **52**:2108-2119.

Sprung, M. 1987. Ecological requirements of developing *Dreissena polymorpha* eggs. Arch. Hydrobiol. Suppl. **79**:69-86.

Sprung, M. 1989. Field and laboratory observations of *Dreissena polymorpha* larvae: abundance, growth, mortality, and food demands. Arch. Hydrobiol. **115**:537-561.

Sprung, M. 1993. The other life: an account of present knowledge of the larval phase of *Dreissena polymorpha*. In: Nalepa, T.F., Scholesser, D.W. (eds) Zebra Mussels: Biology, impacts, and control. Lewis Publishers, CRC Press, Boca Raton, FLA, pp. 39-53.

Sprung, M., and Rose, U. 1988, Influence of food size and food quality on the feeding of the mussel *Dreissena polymorpha*. Oecologia **77**:526-532.

Stańczykowska, A. 1977. Ecology of *Dreissena polymorpha* (Pall) (Bivalvia) in lakes. Pol. Arch. Hydrobiol. **24**:461-530.

Stoeckel, J.A., Schneider, D.W., Soeken, L.A., Blodgett, K.D and Sparks, R.E. 1997. Larval dynamics of a riverine metapopulation: implications for Zebra Mussel recruitment, dispersal, and control in a large-river system. J. NA Benthol. Soc. **16**:586-601.

Stoeckmann, A. 2003. Physiological energetics of Lake Erie dreissenid mussels: a basis for the displacement of *Dreissena polymorpha* by *Dreissena bugensis*. Can. J. Fish. Aquat. Sci. **60**:126-134.

Strayer, D.L., Hattala, K.A., and Kahnle, A.W. 2004. Effects of an invasive bivalve (*Dreissena polymorpha*) on fish in the Hudson River estuary. Can. J. Fish. Aquat. Sci. **61**:924-941.

Therriault, T.W., Docker, M.F., Orlova, M.I., Heath, D.D, and MacIsaac, H.J. 2004. Molecular resolution of the family Dreissenidae (Mollusca: Bivalvia) with emphasis on Ponto-Caspian species, including first report of *Mytilopsis leucophaeata* in the Black Sea basin. Mol. Phylogen. Evol. **30**:479-489.

Therriault, T.W., Orlova, M.I., Docker, M.F., MacIsaac, H.J., and Heath, D.D. 2005. Invasion genetics of a freshwater mussel (*Dreissena rostriformis bugensis*) in Eastern Europe: high gene flow and multiple introductions. Heredity **95**:16-23.

Therriault, T. W., and Herborg, L-M. 2008. A qualitative biological risk assessment for vase tunicate *Ciona intestinalis* in Canadian waters: using expert knowledge. ICES J. Mar. Sci. **65**:781-787.

Therriault, T.W., and Orlova, M.I. 2010. Invasion success within the family Dreissenidae: prerequisites, mechanisms, and perspectives. In: Van der Velde, G., Rajagopal, S., Bij de Vaate, A. 2010. (eds). The Zebra Mussel in Europe. Backhuys Publishers, Leiden/Margraf Publishers, Weikersheim. pp. 59-68.

Thorp, J.H., and Casper, A.F. 2003. Importance of biotic interactions in large rivers: An experiment with planktivorous fish, dreissenid mussels and zooplankton in the St. Lawrence River. Riv. Res. Appl. **19**:265-279.

Van der Velde, G., and Platvoet, D. 2007. Quagga Mussels *Dreissena rostriformis bugensis* (Andrusov, 1897) in the Main River (Germany). Aquat. Invasions **2**:261-264.

Van der Velde, G., Rajagopal and S., Bij de Vaate, A. 2010. (eds). The Zebra Mussel in Europe. Backhuys Publishers, Leiden/Margraf Publishers, Weikersheim.

Verween, A., Vincx, M., Mees, J., and Degraer, S. 2005. Seasonal variability of *Mytilopsis leucophaeta* larvae in the harbour of Antwerp: implications for ecologically and economically sound biofouling control. Belg. J. Zool. **135**:91-93.

Verween, A., Vincx, M., and Degraer, S. 2010. *Mytilopsis leucophaeata*: The brackish water equivalent of *Dreissena polymorpha*? A review. In: Van der Velde, G., Rajagopal, S., Bij de Vaate, A. (eds). The Zebra Mussel in Europe. Backhuys Publishers, Leiden/Margraf Publishers, Weikersheim. pp. 29-44.

Voroshilova, I.S., Artamonova, V.S., Makhrov, A.A., and Slyn'ko, Y.V. 2010. Natural Hybridization of Two Mussel Species *Dreissena polymorpha* (Pallas, 1771) and *Dreissena bugensis* (Andrusov, 1897). Biol. Bull. **37**:542–547.

Wacker, A., and Von Elert, E. 2003. Settlement pattern of the Zebra Mussel, *Dreissena polymorpha*, as a function of depth in Lake Constance. Arch. Hydrobiol. **158**:289-301.

Walton, W.C. 1996. Occurrence of Zebra Mussel (*Dreissena polymorpha*) in the oligohaline Hudson River, New York. Estuaries **19**:612-618.

Whittier, T.R., Ringold, R.L., Herlihy, A.T., and Pierson, S.M. 2008. A calcium-based invasion risk assessment for Zebra and Quagga Mussels (*Dreissena* spp). Front. Ecol. Environ. **6**:180-184.

Wilson, A.B., Boulding, E.G., and Naish, K.A. 1999. Characterization of tri- and tetranucleotide microsatellite loci in the invasive mollusc *Dreissena bugensis*. Mol. Ecol. **8**:692-693.

Wolff, W.J. 1969. The mollusca of the estuarine region of the rivers Rhine, Meuse and Scheldt in relation to the hydrography of the area. II. The Dreissenidae. Basteria **33**:93–103

Wong, W.H., Levinton, J.S., Twining, B.S., and Fisher, N.S. 2003. Assimilation of micro-and mesozooplankton by Zebra Mussels: A demonstration of the food web link between zooplankton and benthic suspension feeders. Limnol. Oceanogr. **48**:308-312.

Zhu, B., Fitzgerald, D.G., Mayer, C.M., Rudstam, L.G., and Mills, E.L. 2006. Alteration of ecosystem function by Zebra Mussels in Oneida Lake: impacts on submerged macrophytes. Ecosystems **9**:1017-1028.

# TABLES

*Table 1. Calcium suitability for Zebra Mussel (Dreissena polymorpha) and Quagga Mussel (Dreissena rostriformis bugensis) based on literature accounts (Cohen and Weinstein, 2001; Cohen et al., 2001; Cohen, 2007; Mackie and Claudi, 2010; Whittier et al., 2008; Benson et al., 2012a; 2012b).*

| Category | Definition | Zebra Mussel Ca (mg/L) | Quagga Mussel Ca (mg/L) |
|---|---|---|---|
| Very Low | No adult survival | < 12 | < 12 |
| Moderate | Evidence that both adult survival AND reproduction are supported at a minimum level | 12 -19 | N/A |
| High | Evidence that good sized populations are supported in terms of both survival and reproduction | 20 - 25 | 12 – 32 |
| Very High | Very close to or at optimal range for all stages of the mussel life history; usually supports high to very high level of infestation | > 25 | > 32 |

*Table 2. Location and earliest known date of world-wide introductions, in chronological order, of the Zebra Mussel, Dreissena polymorpha.*

| Location | Date | Established | Reference |
|---|---|---|---|
| Russia | 1769 | Y | Ludyanskiy *et al.* 1993 |
| Caspian Sea | 1771 | Y | Ludyanskiy *et al.* 1993 |
| Hungary | 1794 | Y | Minchin *et al.* 2002 |
| Lithuania | 1803 | Y | Lithuanian Invasive Species Database 2005 |
| England (London) | 1824 | Y | Kerney and Morton 1970 |
| Netherlands | 1826 | Y | Kerney and Morton 1970 |
| Germany | 1830 | Y | Kerney and Morton 1970 |
| Scotland (Edinburgh) | 1834 | Y | Kerney and Morton 1970 |
| Belgium | 1835 | Y | Belgian Biodiversity Platform 2005 |
| France | 1835 | Y | Kinzelbach 1992 |
| Denmark | 1840 | Y | Kerney and Morton 1970 |
| Estonia | 1840 | Y | Minchin *et al.* 2002 |
| Switzerland | 1860s | Y | Jantz and Schöll 1998 |
| Sweden | 1924 | Y | Danish Forest and Nature Agency 2005 |
| Scandinavia | 1940s | Y | Ludyanskiy *et al.* 1993 |
| Italy | 1969 | Y | Annoni *et al.* 1978 |
| Yugoslavia | 1970s | Y | Ludyanskiy *et al.* 1993 |
| Baltic Sea | 1980s | Y | Orlova *et al.* 2000 |
| Canada (ON) | 1986 | Y | Carlton 2008 |
| United States | 1986 | Y | Carlton 2008 |
| Latvia | 1996 | Y | Minchin *et al.* 2002 |
| Ireland | 1997 | Y | McCarthy *et al.* 1997 |
| Spain | 2001 | Y | Araujo and Álvarez Halcón 2001 |

*Table 3. Ecological impacts associated with Zebra Mussel and Quagga Mussel invasions as reported in the scientific literature (modified from Higgins and VanderZanden, 2010).*

| Element | Survey/Literature Results | | |
|---|---|---|---|
| | Direction | Magnitude | Uncertainty |
| **Physical habitat** | | | |
| Water clarity | Increase | High | Very Low |
| Thermocline depth | Increase | Low | High |
| Littoral zone depth | Increase | Moderate | Low |
| Hard substrate fouling | Increase | High | Very Low |
| Soft substrate fouling | Increase | Moderate | Very Low |
| Deepwater anoxia | Increase | Low | Very High |
| Sediment anoxia | Increase | Moderate | High |
| **Chemical habitat** | | | |
| Particulate nutrients | Decrease | Moderate | Very Low |
| Soluble nutrients (Lakes) | Increase | Low | Very Low |
| Soluble nutrients (Rivers) | Increase | High | Very Low |
| Suspended sediments | Decrease | High | Very Low |
| **Biota** | | | |
| Sediment bacteria | Increase | High | Very Low |
| Phytoplankton (total) | Decrease | High | Very Low |
| Phytoplankton (toxin producing cyanobacteria) | Increase | Moderate | Very Low |
| Periphyton | Increase | High | Very Low |
| Macrophyte cover | Increase | Moderate | Very Low |
| Zooplankton | Decrease | Moderate | Very Low |
| Zoobenthos (littoral) | Increase | High | Very Low |
| Zoobenthos (profundal) | Decrease | Moderate | Very Low |
| Unionid mussel (abundance) | Decrease | Very high | Very Low |
| Fish (planktivore) | Decrease | Moderate | Moderate |
| Fish (benthivore-littoral) | Increase | Moderate | Very Low |
| Fish (deepwater benthivore) | Decrease | High | Moderate |
| Fish (piscivore) | Decrease | Moderate | Moderate |
| Avian botulism | Increase | Moderate | High |
| **Biodiversity** | | | |
| Unionid mussel | Decrease | Very High | Very Low |
| Sphaeriid mussel | Decrease | Very High | Low |
| Species at Risk | Decrease | Low to High | Very High |
| | | **Impact** | **Uncertainty** |
| **Impact to Environment** | | Very High Negative | Very Low |

*Table 4. Canadian freshwater molluscs designated by COSEWIC.*

| Scientific Name | Common Name | COSEWIC Designation | Range | Note |
|---|---|---|---|---|
| *Villosa fabalis* | Rayed Bean | Endangered | ON | |
| *Villosa iris* | Rainbow Mussel | Endangered | ON | |
| *Lampsilis fasciola* | Wavy-rayed Lampmussel | Endangered | ON | Reduced to Special Concern; Potential overlap with dreissenids |
| *Simpsonais ambigua* | Mudpuppy Mussel | Endangered | ON | |
| *Epioblasma torulosa rangiana* | Northern Riffleshell | Endangered | ON | |
| *Epioblasma triquetra* | Snuffbox | Endangered | ON | |
| *Truncilla donaciformis* | Fawnsfoot | Endangered | ON | |
| *Obovaria olivaria* | Hickorynut | Endangered | ON, QC | |
| *Obovaria subrotunda* | Round Hickorynut | Endangered | ON | |
| *Ptychobranchus fasciolaris* | Kidneyshell | Endangered | ON | |
| *Quadrula quadrula* | Mapleleaf Mussel | Endangered | MB | |
| *Gonidea angulata* | Rocky Mountain Ridged Mussel | Endangered | BC | Potential threat from dreissenids |
| *Pleurobema sintoxia* | Round Pigtoe | Endangered | ON | Potential overlap with dreissenids |
| *Ligumia nasuta* | Eastern Pondmussel | Endangered | | Potential overlap with dreissenids |
| *Physella johnsoni* | Banff Springs Snail | Endangered | AB | |
| *Quadrula quadrula* | Mapleleaf Mussel | Threatened | ON | Potential overlap with dreissenids |
| *Alasmidonta varicosa* | Brook Floater | Special Concern | NB, NS | |
| *Lampsilis cariosa* | Yellow Lampmussel | Special Concern | NB, NS | Potential overlap with dreissenids |
| *Acroloxus coloradensis* | Rocky Mountain Capshell | Data Deficient | ON, QC | |
| *Lyrogyrus granum* | Squat Duskysnail | Data Deficient | NB, NS | |
| *Physella parkeri latchfordi* | Gatineau Tadpole Snail | Data Deficient | QC | |

*Table 5. Canadian freshwater fishes designated Extirpated, Endangered, Threatened, or Special Concern by COSEWIC that could potentially be affected by dreissenid mussels.*

| Scientific Name | Common Name | COSEWIC Designation | Range |
|---|---|---|---|
| *Polyodon spathula* | Paddlefish | Extirpated | ON |
| Coregonus sp. | Spring cisco | Endangered | QC |
| *Moxostoma hubbsi* | Copper redhorse | Endangered | QC |
| *Acipenser fulvescens* | Lake sturgeon (various populations) | Endangered | ON, MB, SK, AB, |
| *Lampetra richardsoni* | Western brook lamprey | Endangered | BC |
| *Hybognathus argyritis* | Western silvery minnow | Endangered | AB |
| *Gasterosteus aculeatus* | Enos Lake benthic threespine stickleback | Endangered | BC |
| *Coregonus huntsmani* | Atlantic Whitefish | Endangered | NS |
| *Coregonus zenithicus* | Shortjaw cisco | Threatened | NT, AB, SK, MB, ON |
| *Moxostoma duquesnei* | Black redhorse | Threatened | ON |
| *Rhinichthys Umatilla* | Umatilla dace | Threatened | BC |
| *Lampetra macrostoma* | Vancouver lamprey | Threatened | BC |
| *Cottus aleuticus* | Coastrange sculpin | Threatened | BC |
| *Osmerus mordax* | Rainbow smelt (small-bodied) | Threatened | NB |
| *Catostomus platyrhynchus* | Mountain sucker | Threatened | AB, SK |
| *Acipenser fulvescens* | Lake Sturgeon | Threatened | ON, QC |
| *Ichthyomyzon fossor* | Northern brook lamprey | Special Concern | ON, QC |
| *Ichthyomyzon unicuspis* | Silver lamprey | Special Concern | ON, QC |
| *Moxostoma carinatum* | River redhorse | Special Concern | ON, QC |
| *Myoxocephalus thompsonii* | Deepwater sculpin | Special Concern | ON, QC |
| *Minytrema melanops* | Spotted sucker | Special Concern | ON |
| *Lepomis gulosus* | Warmouth | Special Concern | ON |
| *Ictiobus cyprinellus* | Bigmouth buffalo | Special Concern | ON |
| *Cottus hubbsi* | Columbia sculpin | Special Concern | BC |
| *Cottus confuses* | Shorthead sculpin | Special Concern | BC |
| *Acipenser brevirostrum* | Shortnose sturgeon | Special Concern | NB |
| *Acipenser fulvescens* | Lake sturgeon (various populations) | Special Concern | ON, MB, QC |
| *Catostomus platyrhynchus* | Mountain sucker | Special Concern | BC |

*Table 6. Introduced populations of Mytilopsis leucopheata with reported date and indication of population establishment.*

| Location | Date Reported | Established | Reference |
|---|---|---|---|
| North Sea | 2004 | n/a | Therriault et al., 2004 |
| Black Sea | 2001<br>by 2004 invaded Dniester Liman | yes | Therriault et al., 2004; NOBANIS 2011 |
| Baltic Sea | 2000 | n/a | NOBANIS, 2011 |
| Caspian Sea, Russia | 2004 | n/a | Therriault et al., 2004 |
| Sea of Azov | 2004 | n/a | Therriault et al., 2004 |
| Belgium | 1835 | yes | NOBANIS, 2011; Laine, et al., 2000 |
| Brazil | 2010 | n/a | Kennedy, 2010 |
| Central Gulf of Finland | 2003 | yes | Laine et al., 2006 |
| France | 1898 | n/a | Rajagopal et al., 2005b |
| Germany | 1932<br>1928 sighting in the Kiel Canal | n/a | Rajagopal et al., 2005b; Boettger, 1933 in Verween et al., 2010 |
| Netherlands | 2002<br>by 1969 invaded Rhine River | n/a | Wolff, 1969; Rajagopal et al., 2002b |
| United Kingdom | 2004 | n/a | Therriault et al., 2004 |
| Wales, UK | 1996 | n/a | NOBANIS, 2011 |
| Spain | 2003 | n/a | Escot et al. 2003, in Verween et al., 2010 |
| United States | | | |
| Upper Mississippi River | 1988 | n/a | Koch, 1989; Therriault et al., 2004 |
| Housatonic River, Connecticut | 2010 | yes | Kennedy, 2010 |
| Charles River, Massachusetts | 2010 | yes | Kennedy, 2010 |
| Chesapeake Bay | 1934 | n/a | Johnson, 1934 |
| Chesapeake Bay | 2006 | yes | Verween et al., 2006; NOBANIS, 2011 |
| Several locations in Florida | 2010 | yes | Kennedy, 2010 |
| Several sites in Hudson River | 1992 | yes | Walton, 1996; Verween et al., 2010 |
| New England | 1996 | n/a | Smith and Boss, 1996 |

*Table 7. List of main data sources used to calculate 75$^{th}$ percentile calcium concentrations for sub-drainages used in the risk assessment.*

| Province | Number of sites | Year | Source | Contact |
|---|---|---|---|---|
| British Columbia | 3545 | 1969-2011 | Government of British Columbia | L.-M. Herborg |
| Alberta | 478 | 1981-2009 | Alberta Environment | M. Raven |
| Manitoba | 1145 | 1973-2011 | Manitoba Water Stewardship | J. Shead |
| Ontario* | 8882 | 1970s-2009 | 1970s Lake Inventory Database, Ontario Ministry of Natural Resources; Hincks and Mackie 1997; Beeton et al. 1967; STAR database; Cohen and Weinstein 2001 (pers. comm. therein) | G. Mackie, K. Minns |
| Quebec | 3137 | 1999-2000 | Ministère des Ressources naturelles et de la Faune | A. Paquet, A. Simard |
| Saskatchewan | 119 | 1976-2010 | Government of Saskatchewan | T. Johnston |

*Calcium concentrations were derived from alkalinity using the following relationship: Calcium (mg/L) = Alkalinity (mg $CaCO_3$/L)/3.49 (Mackie and Claudi, 2010).

*Table 8. Thresholds for propagule pressure and proximity to invaded habitats for Zebra Mussel (Dreissena polymorpha) and Quagga Mussel (Dreissena rostriformis bugensis).*

| Probability of Arrival | | | | |
|---|---|---|---|---|
| Propagule Pressure | | | Correction for Proximity | |
| Category | Human footprint index | Vector Pressure | Proximity to Invaded Habitats | Proximity correction factor |
| Very Low | 0 – 3 | 1 | > 2 watersheds from infestation | 0 |
| Low | 4 – 8 | 2 | Invaded, adjacent or 2 sub-drainages away | +1 |
| Moderate | 9 – 19 | 3 | | |
| High | 20 – 31 | 4 | | |
| Very High | 32 – 52 | 5 | | |

*Table 9. Definition of level of impact and categories for the dreissenid mussel risk assessment (modified from Therriault and Herborg, 2008).*

| Impacts | |
|---|---|
| **Category** | **Definition** |
| Very Low Negative | No measurable impact; consequences can be absorbed without additional management action. |
| Low Negative | A measurable limited impact; disruption to the factor in question but reversible or limited in time, space or severity. |
| Moderate Negative | A measurable widespread impact; widespread disruption to the factor in question but reversible or of limited severity or duration. |
| High Negative | A significant impact; widespread disruption to the factor in question that persists over time or is likely not reversible. |
| Very High Negative | A critical impact; extensive disruption to the factor in question that is irreversible. |

| Uncertainty | |
|---|---|
| **Category** | **Definition** |
| Very High | Little or no information; opinion based on general species knowledge. |
| High | Limited information; third party observational evidence or based on circumstantial evidence. |
| Moderate | Moderate level of information; first hand knowledge and/or unsystematic observations. |
| Low | Substantial scientific information; non peer-reviewed information. |
| Very Low | Extensive scientific information; peer-reviewed information. |

*Table 10. Percentage of sites falling into each calcium category for "n" sites and scores for the probability of survival (habitat suitability) for Zebra Mussel per sub-drainage based on calcium concentrations (mg/L; 75$^{th}$ percentile) and corrected for temperature. Sub-drainages were ranked on their suitability for Zebra Mussel survival based on literature accounts (see Table 1).*

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-19 | 20-25 | ≥ 26 | Probability of Survival |
|---|---|---|---|---|---|---|---|---|
| AB | 05F | Battle | 15 | 20% | 27% | 7% | 47% | Very high |
| AB | 06A | Beaver (Alta.-Sask.) | 47 | 6% | 13% | 19% | 62% | Very high |
| AB | 05B | Bow | 21 | 10% | 0% | 0% | 90% | Very high |
| AB | 07C | Central Athabasca – Lower | 19 | 21% | 32% | 11% | 37% | Very high |
| AB | 07B | Central Athabasca – Upper | 47 | 11% | 17% | 21% | 51% | Very high |
| AB | 05E | Central North Saskatchewan | 45 | 11% | 16% | 24% | 49% | Very high |
| AB | 07J | Central Peace – Lower | 28 | 36% | 18% | 18% | 29% | Very high |
| AB | 07H | Central Peace – Upper | 21 | 10% | 19% | 14% | 57% | Very high |
| AB | 08N | Columbia - U.S.A. | 747 | 19% | 17% | 14% | 50% | Very high |
| AB | 10C | Fort Nelson | 7 | 29% | 0% | 0% | 71% | Very high |
| AB | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | 0% | Very low |
| AB | 07O | Hay | 13 | 23% | 15% | 15% | 46% | Very high |
| AB | 71 | Lake Athabasca | 2 | 50% | 0% | 0% | 50% | High |
| AB | 07M | Lake Athabasca – Shores | 6 | 100% | 0% | 0% | 0% | Very low |
| AB | 07D | Lower Athabasca | 37 | 41% | 35% | 11% | 14% | Moderate |
| AB | 05G | Lower North Saskatchewan | 5 | 40% | 20% | 20% | 20% | High |
| AB | 07K | Lower Peace | 15 | 87% | 7% | 0% | 7% | Very low |
| AB | 05H | Lower South Saskatchewan | 42 | 0% | 0% | 0% | 100% | Very high |
| AB | 11A | Missouri | 3 | 0% | 33% | 0% | 67% | Very high |
| AB | 05C | Red Deer | 35 | 14% | 11% | 3% | 71% | Very high |
| AB | 07N | Slave | 10 | 100% | 0% | 0% | 0% | Very low |
| AB | 07G | Smoky | 33 | 3% | 18% | 27% | 52% | Very high |
| AB | 07P | Southern Great Slave Lake | 17 | 53% | 35% | 6% | 6% | Moderate |
| AB | 07A | Upper Athabasca | 9 | 0% | 33% | 0% | 67% | Very high |
| AB | 06B | Upper Churchill (Man.) | 1 | 0% | 0% | 0% | 100% | Very high |
| AB | 08K | Upper Fraser | 198 | 21% | 22% | 12% | 45% | Very high |
| AB | 05D | Upper North Saskatchewan | 12 | 8% | 8% | 17% | 67% | Very high |
| AB | 07F | Upper Peace | 157 | 3% | 11% | 10% | 76% | Very high |
| AB | 05A | Upper South Saskatchewan | 52 | 0% | 6% | 8% | 87% | Very high |
| BC | 08A | Alsek | 5 | 0% | 0% | 60% | 40% | High |
| BC | 05B | Bow | 21 | 10% | 0% | 0% | 90% | Very high |
| BC | 08F | Central Coastal Waters of B.C. | 88 | 90% | 6% | 0% | 5% | Very low |
| BC | 10B | Central Liard | 14 | 43% | 21% | 0% | 36% | Very high |
| BC | 08N | Columbia - U.S.A. | 747 | 19% | 17% | 14% | 50% | Very high |
| BC | 10C | Fort Nelson | 7 | 29% | 0% | 0% | 71% | Very high |
| BC | 07O | Hay | 13 | 23% | 15% | 15% | 46% | Very high |
| BC | 09A | Headwaters Yukon | 38 | 61% | 26% | 5% | 8% | Very low |
| BC | 08M | Lower Fraser | 299 | 34% | 26% | 14% | 25% | High |
| BC | 08D | Nass – Coast | 64 | 55% | 27% | 13% | 6% | Moderate |
| BC | 08J | Nechako | 204 | 48% | 33% | 8% | 11% | Moderate |
| BC | 08B | Northern Coastal Waters of B.C. | 27 | 19% | 37% | 26% | 19% | Moderate |
| BC | 08O | Queen Charlotte Islands | 30 | 90% | 7% | 0% | 3% | Very low |
| BC | 08E | Skeena – Coast | 309 | 57% | 25% | 6% | 12% | Moderate |
| BC | 07G | Smoky | 33 | 3% | 18% | 27% | 52% | Very high |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-19 | 20-25 | ≥ 26 | Probability of Survival |
|---|---|---|---|---|---|---|---|---|
| BC | 08G | Southern Coastal Waters of B.C. | 92 | 80% | 11% | 2% | 7% | Very low |
| BC | 08C | Stikine – Coast | 62 | 40% | 18% | 19% | 23% | Moderate |
| BC | 08L | Thompson | 520 | 20% | 21% | 10% | 49% | Very high |
| BC | 07A | Upper Athabasca | 9 | 0% | 33% | 0% | 67% | Very high |
| BC | 08K | Upper Fraser | 198 | 21% | 22% | 12% | 45% | Very high |
| BC | 10A | Upper Liard | 23 | 30% | 17% | 9% | 43% | High |
| BC | 05D | Upper North Saskatchewan | 12 | 8% | 8% | 17% | 67% | Very high |
| BC | 07F | Upper Peace | 157 | 3% | 11% | 10% | 76% | Very high |
| BC | 05A | Upper South Saskatchewan | 52 | 0% | 6% | 8% | 87% | Very high |
| BC | 08H | Vancouver Island | 562 | 82% | 10% | 2% | 5% | Very low |
| BC | 07E | Williston Lake | 68 | 19% | 38% | 10% | 32% | Very high |
| MB | 05M | Assiniboine | 135 | 0% | 3% | 4% | 93% | Very high |
| MB | 06E | Central Churchill (Man.) – Lower | 68 | 50% | 16% | 1% | 32% | Very high |
| MB | 05R | Eastern Lake Winnipeg | 130 | 86% | 9% | 1% | 4% | Very low |
| MB | 05T | Grass and Burntwood | 76 | 8% | 68% | 9% | 14% | Moderate |
| MB | 04A | Hayes (Man.) | 17 | 12% | 53% | 18% | 18% | High |
| MB | 50 | Lake Winnipeg | 99 | 5% | 9% | 14% | 72% | Very high |
| MB | 05L | Lake Winnipegosis and Lake Manitoba | 105 | 0% | 1% | 1% | 98% | Very high |
| MB | 06F | Lower Churchill (Man.) | 16 | 0% | 38% | 50% | 13% | High |
| MB | 05U | Nelson | 39 | 0% | 10% | 23% | 67% | Very high |
| MB | 05J | Qu'Appelle | 40 | 0% | 0% | 0% | 100% | Very high |
| MB | 05O | Red | 150 | 0% | 2% | 1% | 97% | Very high |
| MB | 06D | Reindeer | 1 | 100% | 0% | 0% | 0% | Very low |
| MB | 05K | Saskatchewan | 156 | 0% | 13% | 9% | 78% | Very high |
| MB | 04C | Severn | 77 | 43% | 49% | 8% | 0% | Moderate |
| MB | 05N | Souris | 35 | 0% | 0% | 3% | 97% | Very high |
| MB | 05S | Western Lake Winnipeg | 65 | 2% | 5% | 3% | 91% | Very high |
| MB | 05P | Winnipeg | 955 | 81% | 11% | 4% | 4% | Very low |
| NL | 03L | Caniapiscau | 10 | 100% | 0% | 0% | 0% | Very low |
| NL | 03O | Churchill (Nfld.) | 1 | 100% | 0% | 0% | 0% | Very low |
| NL | 03M | Eastern Ungava Bay | 78 | 100% | 0% | 0% | 0% | Very low |
| NL | 02W | Gulf of St. Lawrence – Natashquan | 42 | 93% | 2% | 0% | 5% | Very low |
| NL | 02V | Gulf of St. Lawrence – Romaine | 39 | 100% | 0% | 0% | 0% | Very low |
| NL | 02U | Moisie and St. Lawrence Estuary | 61 | 100% | 0% | 0% | 0% | Very low |
| NL | 02X | Petit Mécatina and Strait of Belle Isle | 5 | 100% | 0% | 0% | 0% | Very low |
| NT | 10B | Central Liard | 14 | 43% | 21% | 0% | 36% | Very high |
| NT | 07L | Fond-du-Lac | 1 | 100% | 0% | 0% | 0% | Very low |
| NT | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | 0% | Very low |
| NT | 07O | Hay | 13 | 23% | 15% | 15% | 46% | Very high |
| NT | 07N | Slave | 10 | 100% | 0% | 0% | 0% | Very low |
| NT | 07P | Southern Great Slave Lake | 17 | 53% | 35% | 6% | 6% | Moderate |
| NT | 10A | Upper Liard | 23 | 30% | 17% | 9% | 43% | High |
| ON | 04M | Abitibi | 388 | 44% | 28% | 10% | 18% | High |
| ON | 04F | Attawapiskat – Coast | 46 | 57% | 33% | 2% | 9% | Moderate |
| ON | 02K | Central Ottawa | 1098 | 70% | 11% | 5% | 13% | Moderate |
| ON | 02E | Eastern Georgian Bay | 752 | 91% | 2% | 1% | 6% | Very low |
| ON | 02F | Eastern Lake Huron | 186 | 5% | 3% | 3% | 88% | Very high |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-19 | 20-25 | ≥ 26 | Probability of Survival |
|---|---|---|---|---|---|---|---|---|
| ON | 05R | Eastern Lake Winnipeg | 130 | 86% | 9% | 1% | 4% | Very low |
| ON | 04E | Ekwan – Coast | 10 | 0% | 60% | 20% | 20% | High |
| ON | 05Q | English | 765 | 86% | 12% | 2% | 1% | Very low |
| ON | 20 | Great Lakes and St. Lawrence | 39 | 38% | 18% | 8% | 36% | Very high |
| ON | 04N | Harricanaw – Coast | 77 | 91% | 6% | 0% | 3% | Very low |
| ON | 04A | Hayes (Man.) | 17 | 12% | 53% | 18% | 18% | High |
| ON | 04J | Kenogami | 314 | 13% | 25% | 21% | 41% | Very high |
| ON | 02H | Lake Ontario and Niagara Peninsula | 539 | 55% | 12% | 7% | 25% | High |
| ON | 04H | Lower Albany – Coast | 3 | 100% | 0% | 0% | 0% | Very low |
| ON | 02L | Lower Ottawa | 503 | 82% | 9% | 3% | 7% | Very low |
| ON | 04L | Missinaibi-Mattagami | 646 | 51% | 25% | 11% | 13% | Moderate |
| ON | 04K | Moose (Ont.) | 16 | 50% | 44% | 0% | 6% | Moderate |
| ON | 02B | Northeastern Lake Superior | 649 | 70% | 18% | 6% | 6% | Moderate |
| ON | 02G | Northern Lake Erie | 95 | 2% | 3% | 4% | 91% | Very high |
| ON | 02C | Northern Lake Huron | 850 | 96% | 2% | 0% | 1% | Very low |
| ON | 02A | Northwestern Lake Superior | 347 | 64% | 23% | 7% | 6% | Moderate |
| ON | 04C | Severn | 77 | 43% | 49% | 8% | 0% | Moderate |
| ON | 04G | Upper Albany | 229 | 72% | 21% | 5% | 3% | Moderate |
| ON | 02J | Upper Ottawa | 854 | 89% | 7% | 2% | 3% | Very low |
| ON | 02M | Upper St. Lawrence | 89 | 25% | 21% | 17% | 37% | Very high |
| ON | 02D | Wanipitai and French (Ont.) | 374 | 94% | 4% | 1% | 0% | Very low |
| ON | 04D | Winisk – Coast | 45 | 42% | 42% | 11% | 4% | Moderate |
| ON | 05P | Winnipeg | 955 | 81% | 11% | 4% | 4% | Very low |
| QC | 04M | Abitibi | 388 | 44% | 28% | 10% | 18% | High |
| QC | 02S | Betsiamites – Coast | 105 | 100% | 0% | 0% | 0% | Very low |
| QC | 03B | Broadback and Rupert | 92 | 90% | 7% | 0% | 3% | Very low |
| QC | 03L | Caniapiscau | 10 | 100% | 0% | 0% | 0% | Very low |
| QC | 02K | Central Ottawa | 1098 | 70% | 11% | 5% | 13% | Moderate |
| QC | 02O | Central St. Lawrence | 251 | 61% | 21% | 5% | 14% | Moderate |
| QC | 03O | Churchill (Nfld.) | 1 | 100% | 0% | 0% | 0% | Very low |
| QC | 03M | Eastern Ungava Bay | 78 | 100% | 0% | 0% | 0% | Very low |
| QC | 03C | Eastmain | 6 | 100% | 0% | 0% | 0% | Very low |
| QC | 20 | Great Lakes and St. Lawrence | 39 | 38% | 18% | 8% | 36% | Very high |
| QC | 02W | Gulf of St. Lawrence – Natashquan | 42 | 93% | 2% | 0% | 5% | Very low |
| QC | 02V | Gulf of St. Lawrence – Romaine | 39 | 100% | 0% | 0% | 0% | Very low |
| QC | 01B | Gulf of St. Lawrence and Northern Bay of Fundy (N.B.) | 37 | 19% | 19% | 19% | 43% | Very high |
| QC | 04N | Harricanaw – Coast | 77 | 91% | 6% | 0% | 3% | Very low |
| QC | 03D | La Grande – Coast | 1 | 100% | 0% | 0% | 0% | Very low |
| QC | 02L | Lower Ottawa | 503 | 82% | 9% | 3% | 7% | Very low |
| QC | 02P | Lower St. Lawrence | 385 | 95% | 3% | 1% | 1% | Very low |
| QC | 02T | Manicouagan and aux Outardes | 92 | 100% | 0% | 0% | 0% | Very low |
| QC | 02U | Moisie and St. Lawrence Estuary | 61 | 100% | 0% | 0% | 0% | Very low |
| QC | 02Q | Northern Gaspé Peninsula | 41 | 20% | 27% | 22% | 32% | Very high |
| QC | 03A | Nottaway – Coast | 209 | 97% | 3% | 0% | 0% | Very low |
| QC | 02X | Petit Mécatina and Strait of Belle Isle | 5 | 100% | 0% | 0% | 0% | Very low |
| QC | 02R | Saguenay | 343 | 99% | 0% | 0% | 0% | Very low |
| QC | 01A | Saint John and Southern Bay of | 35 | 34% | 34% | 17% | 14% | High |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-19 | 20-25 | ≥ 26 | Probability of Survival |
|---|---|---|---|---|---|---|---|---|
| | | Fundy (N.B.) | | | | | | |
| QC | 02N | Saint-Maurice | 313 | 100% | 0% | 0% | 0% | Very low |
| QC | 02J | Upper Ottawa | 854 | 89% | 7% | 2% | 3% | Very low |
| QC | 02M | Upper St. Lawrence | 89 | 25% | 21% | 17% | 37% | Very high |
| SK | 05M | Assiniboine | 135 | 0% | 3% | 4% | 93% | Very high |
| SK | 05F | Battle | 15 | 20% | 27% | 7% | 47% | Very high |
| SK | 06A | Beaver (Alta.-Sask.) | 47 | 6% | 13% | 19% | 62% | Very high |
| SK | 07C | Central Athabasca – Lower | 19 | 21% | 32% | 11% | 37% | Very high |
| SK | 06E | Central Churchill (Man.) – Lower | 68 | 50% | 16% | 1% | 32% | Very high |
| SK | 06C | Central Churchill (Man.) – Upper | 1 | 0% | 100% | 0% | 0% | Moderate |
| SK | 05E | Central North Saskatchewan | 45 | 11% | 16% | 24% | 49% | Very high |
| SK | 07L | Fond-du-Lac | 1 | 100% | 0% | 0% | 0% | Very low |
| SK | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | 0% | Very low |
| SK | 71 | Lake Athabasca | 2 | 50% | 0% | 0% | 50% | High |
| SK | 07M | Lake Athabasca – Shores | 6 | 100% | 0% | 0% | 0% | Very low |
| SK | 05L | Lake Winnipegosis and Lake Manitoba | 105 | 0% | 1% | 1% | 98% | Very high |
| SK | 07D | Lower Athabasca | 37 | 41% | 35% | 11% | 14% | Moderate |
| SK | 05G | Lower North Saskatchewan | 5 | 40% | 20% | 20% | 20% | High |
| SK | 05H | Lower South Saskatchewan | 42 | 0% | 0% | 0% | 100% | Very high |
| SK | 11A | Missouri | 3 | 0% | 33% | 0% | 67% | Very high |
| SK | 05J | Qu'Appelle | 40 | 0% | 0% | 0% | 100% | Very high |
| SK | 05C | Red Deer | 35 | 14% | 11% | 3% | 71% | Very high |
| SK | 06D | Reindeer | 1 | 100% | 0% | 0% | 0% | Very low |
| SK | 05K | Saskatchewan | 156 | 0% | 13% | 9% | 78% | Very high |
| SK | 05N | Souris | 35 | 0% | 0% | 3% | 97% | Very high |
| SK | 06B | Upper Churchill (Man.) | 1 | 0% | 0% | 0% | 100% | Very high |
| SK | 05A | Upper South Saskatchewan | 52 | 0% | 6% | 8% | 87% | Very high |
| YT | 08A | Alsek | 5 | 0% | 0% | 60% | 40% | High |
| YT | 10B | Central Liard | 14 | 43% | 21% | 0% | 36% | Very high |
| YT | 09A | Headwaters Yukon | 38 | 61% | 26% | 5% | 8% | Very low |
| YT | 10A | Upper Liard | 23 | 30% | 17% | 9% | 43% | High |

*Table 11. Probability of Zebra Mussel arrival, survival, and invasion per sub-drainage. The probability of invasion is based on the probability of survival (calcium suitability corrected for temperature) and probability of arrival (propagule pressure corrected for proximity to an invaded watershed). The risk to the environment is based on the probability of invasion and impacts to the environment.*

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| AB | 05F | Battle | Very high | 0 | Very high | High | 0 | High | Very high | High |
| AB | 06A | Beaver (Alta.-Sask.) | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 05B | Bow | Very high | 0 | Very high | High | 0 | High | Very high | High |
| AB | 07C | Central Athabasca - Lower | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 07B | Central Athabasca - Upper | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 05E | Central North Saskatchewan | Very high | 0 | Very high | High | 0 | High | Very high | High |
| AB | 07J | Central Peace - Lower | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 07H | Central Peace - Upper | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 08N | Columbia - U.S.A. | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 10C | Fort Nelson | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 07Q | Great Slave Lake - East Arm South Shore | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 07O | Hay | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 71 | Lake Athabasca | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| AB | 07M | Lake Athabasca - Shores | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 07D | Lower Athabasca | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| AB | 05G | Lower North Saskatchewan | High | 0 | High | High | 0 | High | High | High |
| AB | 07K | Lower Peace | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 05H | Lower South Saskatchewan | Very high | 0 | Very high | High | 0 | High | Very high | High |
| AB | 11A | Missouri | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| AB | 05C | Red Deer | Very high | 0 | Very high | High | 0 | High | Very high | High |
| AB | 07N | Slave | Very low | 0 | Very low | Low | 0 | Low | Very low | Low |
| AB | 07G | Smoky | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 07P | Southern Great Slave Lake | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| AB | 07A | Upper Athabasca | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 06B | Upper Churchill (Man.) | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| AB | 08K | Upper Fraser | Very high | 0 | Very high | Low | 0 | Low | High | High |
| AB | 05D | Upper North Saskatchewan | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 07F | Upper Peace | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| AB | 05A | Upper South Saskatchewan | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08A | Alsek | Very high | -1 | High | Low | 0 | Low | Moderate | High |

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| BC | 05B | Bow | Very high | 0 | Very high | High | 0 | High | Very high | High |
| BC | 08F | Central Coastal Waters of B.C. | Very low | 0 | Very low | Low | 0 | Low | Very low | Low |
| BC | 10B | Central Liard | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| BC | 08N | Columbia - U.S.A. | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 10C | Fort Nelson | Very high | 0 | Very high | Low | 0 | Low | High | High |
| BC | 07O | Hay | Very high | 0 | Very high | Low | 0 | Low | High | High |
| BC | 09A | Headwaters Yukon | Moderate | -1 | Low | Low | 0 | Low | Low | Low |
| BC | 08M | Lower Fraser | High | 0 | High | Moderate | 0 | Moderate | High | High |
| BC | 08D | Nass - Coast | Moderate | 0 | Moderate | Low | 0 | Low | Moderate | High |
| BC | 08J | Nechako | Moderate | 0 | Moderate | Low | 0 | Low | Moderate | High |
| BC | 08B | Northern Coastal Waters of B.C. | High | -1 | Moderate | Low | 0 | Low | Moderate | High |
| BC | 08O | Queen Charlotte Islands | Very low | 0 | Very low | Moderate | 0 | Moderate | Very low | Low |
| BC | 08E | Skeena - Coast | Moderate | 0 | Moderate | Low | 0 | Low | Moderate | High |
| BC | 07G | Smoky | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08G | Southern Coastal Waters of B.C. | Very low | 0 | Very low | Low | 0 | Low | Very low | Low |
| BC | 08C | Stikine - Coast | High | -1 | Moderate | Low | 0 | Low | Moderate | High |
| BC | 08L | Thompson | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 07A | Upper Athabasca | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08K | Upper Fraser | Very high | 0 | Very high | Low | 0 | Low | High | High |
| BC | 10A | Upper Liard | Very high | -1 | High | Low | 0 | Low | Moderate | High |
| BC | 05D | Upper North Saskatchewan | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 07F | Upper Peace | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 05A | Upper South Saskatchewan | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08H | Vancouver Island | Very low | 0 | Very low | Moderate | 0 | Moderate | Very low | Low |
| BC | 07E | Williston Lake | Very high | 0 | Very high | Low | 0 | Low | High | High |
| MB | 05M | Assiniboine | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| MB | 06E | Central Churchill (Man.) - Lower | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| MB | 05R | Eastern Lake Winnipeg | Very low | 0 | Very low | Very low | 1 | Low | Very low | Low |
| MB | 05T | Grass and Burntwood | Moderate | 0 | Moderate | Low | 0 | Low | Moderate | High |
| MB | 04A | Hayes (Man.) | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| MB | 50 | Lake Winnipeg | Very high | 0 | Very high | Low | 1 | Very high | Very high | High |
| MB | 05L | Lake Winnipegosis and Lake | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| | | Manitoba | | | | | | | | |
| MB | 06F | Lower Churchill (Man.) | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| MB | 05U | Nelson | Very high | 0 | Very high | Very low | 1 | Very high | Very high | High |
| MB | 05J | Qu'Appelle | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| MB | 05O | Red | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| MB | 06D | Reindeer | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| MB | 05K | Saskatchewan | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| MB | 04C | Severn | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| MB | 05N | Souris | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| MB | 05S | Western Lake Winnipeg | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| MB | 05P | Winnipeg | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| NL | 03L | Caniapiscau | Very low | -1 | Very low | Very low | 1 | Low | Very low | Low |
| NL | 03O | Churchill (Nfld.) | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 03M | Eastern Ungava Bay | Very low | -1 | Very low | Low | 0 | Low | Very low | Low |
| NL | 02W | Gulf of St. Lawrence - Natashquan | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 02V | Gulf of St. Lawrence - Romaine | Very low | 0 | Very low | Very low | 1 | Low | Very low | Low |
| NL | 02U | Moisie and St. Lawrence Estuary | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 02X | Petit Mécatina and Strait of Belle Isle | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| NT | 10B | Central Liard | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| NT | 07L | Fond-du-Lac | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| NT | 07Q | Great Slave Lake - East Arm South Shore | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| NT | 07O | Hay | Very high | 0 | Very high | Low | 0 | Low | High | High |
| NT | 07N | Slave | Very low | 0 | Very low | Low | 0 | Low | Very low | Low |
| NT | 07P | Southern Great Slave Lake | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| NT | 10A | Upper Liard | Very high | -1 | High | Low | 0 | Low | Moderate | High |
| ON | 04M | Abitibi | High | 0 | High | Moderate | 1 | High | High | High |
| ON | 04F | Attawapiskat - Coast | Moderate | 0 | Moderate | Very low | 1 | Low | Moderate | High |
| ON | 02K | Central Ottawa | Moderate | 0 | Moderate | Moderate | 1 | High | High | High |
| ON | 02E | Eastern Georgian Bay | Very low | 0 | Very low | High | 1 | Very high | Very low | Low |
| ON | 02F | Eastern Lake Huron | Very high | 0 | Very high | Very high | 1 | Very high | Very high | High |
| ON | 05R | Eastern Lake Winnipeg | Very low | 0 | Very low | Very low | 1 | Low | Very low | Low |

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| ON | 04E | Ekwan - Coast | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| ON | 05Q | English | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| ON | 20 | Great Lakes and St. Lawrence | Very high | 0 | Very high | Very high | 1 | Very high | Very high | High |
| ON | 04N | Harricanaw - Coast | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| ON | 04A | Hayes (Man.) | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| ON | 04J | Kenogami | Very high | 0 | Very high | Low | 1 | Moderate | High | High |
| ON | 02H | Lake Ontario and Niagara Peninsula | High | 0 | High | Very high | 1 | Very high | Very high | High |
| ON | 04H | Lower Albany - Coast | Very low | 0 | Very low | Very low | 1 | Low | Very low | Low |
| ON | 02L | Lower Ottawa | Very low | 0 | Very low | High | 1 | Very high | Very low | Low |
| ON | 04L | Missinaibi-Mattagami | Moderate | 0 | Moderate | Moderate | 1 | High | High | High |
| ON | 04K | Moose (Ont.) | Moderate | 0 | Moderate | Very low | 1 | Low | Moderate | High |
| ON | 02B | Northeastern Lake Superior | Moderate | 0 | Moderate | Low | 1 | Moderate | Moderate | High |
| ON | 02G | Northern Lake Erie | Very high | 0 | Very high | Very high | 1 | Very high | Very high | High |
| ON | 02C | Northern Lake Huron | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| ON | 02A | Northwestern Lake Superior | Moderate | 0 | Moderate | Moderate | 1 | High | High | High |
| ON | 04C | Severn | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| ON | 04G | Upper Albany | Moderate | 0 | Moderate | Very low | 1 | Low | Moderate | High |
| ON | 02J | Upper Ottawa | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| ON | 02M | Upper St. Lawrence | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| ON | 02D | Wanipitai and French (Ont.) | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| ON | 04D | Winisk - Coast | Moderate | 0 | Moderate | Very low | 1 | Low | Moderate | High |
| ON | 05P | Winnipeg | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| QC | 04M | Abitibi | High | 0 | High | Moderate | 1 | High | High | High |
| QC | 02S | Betsiamites - Coast | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03B | Broadback and Rupert | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03L | Caniapiscau | Very low | -1 | Very low | Very low | 1 | Low | Very low | Low |
| QC | 02K | Central Ottawa | Moderate | 0 | Moderate | Moderate | 1 | High | High | High |
| QC | 02O | Central St. Lawrence | Moderate | 0 | Moderate | Very high | 1 | Very high | High | High |
| QC | 03O | Churchill (Nfld.) | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03M | Eastern Ungava Bay | Very low | -1 | Very low | Low | 0 | Low | Very low | Low |
| QC | 03C | Eastmain | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 20 | Great Lakes and St. Lawrence | Very high | 0 | Very high | Very high | 1 | Very high | Very high | High |
| QC | 02W | Gulf of St. Lawrence - Natashquan | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| QC | 02V | Gulf of St. Lawrence - Romaine | Very low | 0 | Very low | Very low | 1 | Low | Very low | Low |
| QC | 01B | Gulf of St. Lawrence and Northern Bay of Fundy (N.B.) | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| QC | 04N | Harricanaw - Coast | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03D | La Grande - Coast | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02L | Lower Ottawa | Very low | 0 | Very low | High | 1 | Very high | Very low | Low |
| QC | 02P | Lower St. Lawrence | Very low | 0 | Very low | High | 1 | Very high | Very low | Low |
| QC | 02T | Manicouagan and aux Outardes | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02U | Moisie and St. Lawrence Estuary | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02Q | Northern Gaspé Peninsula | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| QC | 03A | Nottaway - Coast | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02X | Petit Mécatina and Strait of Belle Isle | Very low | 0 | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02R | Saguenay | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| QC | 01A | Saint John and Southern Bay of Fundy (N.B.) | High | 0 | High | High | 1 | Very high | Very high | High |
| QC | 02N | Saint-Maurice | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| QC | 02J | Upper Ottawa | Very low | 0 | Very low | Moderate | 1 | High | Very low | Low |
| QC | 02M | Upper St. Lawrence | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| SK | 05M | Assiniboine | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| SK | 05F | Battle | Very high | 0 | Very high | High | 0 | High | Very high | High |
| SK | 06A | Beaver (Alta.-Sask.) | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| SK | 07C | Central Athabasca - Lower | Very high | 0 | Very high | Low | 0 | Low | High | High |
| SK | 06E | Central Churchill (Man.) - Lower | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| SK | 06C | Central Churchill (Man.) - Upper | Moderate | 0 | Moderate | Low | 0 | Low | Moderate | High |
| SK | 05E | Central North Saskatchewan | Very high | 0 | Very high | High | 0 | High | Very high | High |
| SK | 07L | Fond-du-Lac | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 07Q | Great Slave Lake - East Arm South Shore | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 71 | Lake Athabasca | High | 0 | High | Very low | 0 | Very low | Moderate | High |
| SK | 07M | Lake Athabasca - Shores | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |

| Prov. | ID | Sub-drainage | Calcium Suitability | Temp corr. | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|---|---|
| SK | 05L | Lake Winnipegosis and Lake Manitoba | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| SK | 07D | Lower Athabasca | Moderate | 0 | Moderate | Very low | 0 | Very low | Low | Moderate |
| SK | 05G | Lower North Saskatchewan | High | 0 | High | High | 0 | High | High | High |
| SK | 05H | Lower South Saskatchewan | Very high | 0 | Very high | High | 0 | High | Very high | High |
| SK | 11A | Missouri | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| SK | 05J | Qu'Appelle | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| SK | 05C | Red Deer | Very high | 0 | Very high | High | 0 | High | Very high | High |
| SK | 06D | Reindeer | Very low | 0 | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 05K | Saskatchewan | Very high | 0 | Very high | Moderate | 1 | High | Very high | High |
| SK | 05N | Souris | Very high | 0 | Very high | High | 1 | Very high | Very high | High |
| SK | 06B | Upper Churchill (Man.) | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| SK | 05A | Upper South Saskatchewan | Very high | 0 | Very high | Moderate | 0 | Moderate | High | High |
| YT | 08A | Alsek | Very high | -1 | High | Low | 0 | Low | Moderate | High |
| YT | 10B | Central Liard | Very high | 0 | Very high | Very low | 0 | Very low | Moderate | High |
| YT | 09A | Headwaters Yukon | Moderate | -1 | Low | Low | 0 | Low | Low | Low |
| YT | 10A | Upper Liard | Very high | -1 | High | Low | 0 | Low | Moderate | High |

*Table 12. Percentage of sites falling into each calcium category for "n" sites and scores for probability of survival (habitat suitability) for Quagga Mussel per sub-drainage based on calcium concentrations (mg/L; 75$^{th}$ percentile). Sub-drainages were ranked on their suitability for Quagga Mussel survival based on literature accounts (see Table 1).*

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-32 | >32 | Probability of Survival |
|---|---|---|---|---|---|---|---|
| AB | 05F | Battle | 15 | 20% | 53% | 27% | High |
| AB | 06A | Beaver (Alta.-Sask.) | 47 | 6% | 74% | 19% | High |
| AB | 05B | Bow | 21 | 10% | 24% | 67% | Very high |
| AB | 07C | Central Athabasca - Lower | 19 | 21% | 63% | 16% | High |
| AB | 07B | Central Athabasca - Upper | 47 | 11% | 70% | 19% | High |
| AB | 05E | Central North Saskatchewan | 45 | 11% | 62% | 27% | Very high |
| AB | 07J | Central Peace - Lower | 28 | 36% | 46% | 18% | High |
| AB | 07H | Central Peace - Upper | 21 | 10% | 57% | 33% | Very high |
| AB | 08N | Columbia - U.S.A. | 747 | 19% | 45% | 36% | Very high |
| AB | 10C | Fort Nelson | 7 | 29% | 14% | 57% | Very high |
| AB | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | Very low |
| AB | 07O | Hay | 13 | 23% | 46% | 31% | Very high |
| AB | 71 | Lake Athabasca | 2 | 50% | 50% | 0% | High |
| AB | 07M | Lake Athabasca - Shores | 6 | 100% | 0% | 0% | Very low |
| AB | 07D | Lower Athabasca | 37 | 41% | 51% | 8% | High |
| AB | 05G | Lower North Saskatchewan | 5 | 40% | 40% | 20% | High |
| AB | 07K | Lower Peace | 15 | 87% | 7% | 7% | Very low |
| AB | 05H | Lower South Saskatchewan | 42 | 0% | 0% | 100% | Very high |
| AB | 11A | Missouri | 3 | 0% | 67% | 33% | Very high |
| AB | 05C | Red Deer | 35 | 14% | 37% | 49% | Very high |
| AB | 07N | Slave | 10 | 100% | 0% | 0% | Very low |
| AB | 07G | Smoky | 33 | 3% | 52% | 45% | Very high |
| AB | 07P | Southern Great Slave Lake | 17 | 53% | 47% | 0% | High |
| AB | 07A | Upper Athabasca | 9 | 0% | 44% | 56% | Very high |
| AB | 06B | Upper Churchill (Man.) | 1 | 0% | 100% | 0% | High |
| AB | 08K | Upper Fraser | 198 | 21% | 44% | 35% | Very high |
| AB | 05D | Upper North Saskatchewan | 12 | 8% | 75% | 17% | High |
| AB | 07F | Upper Peace | 157 | 3% | 36% | 61% | Very high |
| AB | 05A | Upper South Saskatchewan | 52 | 0% | 42% | 58% | Very high |
| BC | 08A | Alsek | 5 | 0% | 60% | 40% | Very high |
| BC | 05B | Bow | 21 | 10% | 24% | 67% | Very high |
| BC | 08F | Central Coastal Waters of B.C. | 88 | 90% | 8% | 2% | Very low |
| BC | 10B | Central Liard | 14 | 43% | 36% | 21% | High |
| BC | 08N | Columbia - U.S.A. | 747 | 19% | 45% | 36% | Very high |
| BC | 10C | Fort Nelson | 7 | 29% | 14% | 57% | Very high |
| BC | 07O | Hay | 13 | 23% | 46% | 31% | Very high |
| BC | 09A | Headwaters Yukon | 38 | 61% | 37% | 3% | High |
| BC | 08M | Lower Fraser | 299 | 34% | 52% | 13% | High |
| BC | 08D | Nass - Coast | 64 | 55% | 44% | 2% | High |
| BC | 08J | Nechako | 204 | 48% | 43% | 9% | High |
| BC | 08B | Northern Coastal Waters of B.C. | 27 | 19% | 70% | 11% | High |
| BC | 08O | Queen Charlotte Islands | 30 | 90% | 7% | 3% | Very low |
| BC | 08E | Skeena - Coast | 309 | 57% | 35% | 8% | High |
| BC | 07G | Smoky | 33 | 3% | 52% | 45% | Very high |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-32 | >32 | Probability of Survival |
|---|---|---|---|---|---|---|---|
| BC | 08G | Southern Coastal Waters of B.C. | 92 | 80% | 13% | 7% | Very low |
| BC | 08C | Stikine - Coast | 62 | 40% | 40% | 19% | High |
| BC | 08L | Thompson | 520 | 20% | 40% | 40% | Very high |
| BC | 07A | Upper Athabasca | 9 | 0% | 44% | 56% | Very high |
| BC | 08K | Upper Fraser | 198 | 21% | 44% | 35% | Very high |
| BC | 10A | Upper Liard | 23 | 30% | 48% | 22% | High |
| BC | 05D | Upper North Saskatchewan | 12 | 8% | 75% | 17% | High |
| BC | 07F | Upper Peace | 157 | 3% | 36% | 61% | Very high |
| BC | 05A | Upper South Saskatchewan | 52 | 0% | 42% | 58% | Very high |
| BC | 08H | Vancouver Island | 562 | 82% | 15% | 2% | Very low |
| BC | 07E | Williston Lake | 68 | 19% | 60% | 21% | High |
| MB | 05M | Assiniboine | 135 | 0% | 18% | 82% | Very high |
| MB | 06E | Central Churchill (Man.) - Lower | 68 | 50% | 19% | 31% | Very high |
| MB | 05R | Eastern Lake Winnipeg | 130 | 86% | 11% | 3% | Very low |
| MB | 05T | Grass and Burntwood | 76 | 8% | 82% | 11% | High |
| MB | 04A | Hayes (Man.) | 17 | 12% | 82% | 6% | High |
| MB | 50 | Lake Winnipeg | 99 | 5% | 66% | 29% | Very high |
| MB | 05L | Lake Winnipegosis and Lake Manitoba | 105 | 0% | 7% | 93% | Very high |
| MB | 06F | Lower Churchill (Man.) | 16 | 0% | 94% | 6% | High |
| MB | 05U | Nelson | 39 | 0% | 67% | 33% | Very high |
| MB | 05J | Qu'Appelle | 40 | 0% | 3% | 98% | Very high |
| MB | 05O | Red | 150 | 0% | 5% | 95% | Very high |
| MB | 06D | Reindeer | 1 | 100% | 0% | 0% | Very low |
| MB | 05K | Saskatchewan | 156 | 0% | 48% | 52% | Very high |
| MB | 04C | Severn | 77 | 43% | 57% | 0% | High |
| MB | 05N | Souris | 35 | 0% | 9% | 91% | Very high |
| MB | 05S | Western Lake Winnipeg | 65 | 2% | 22% | 77% | Very high |
| MB | 05P | Winnipeg | 955 | 81% | 17% | 2% | Very low |
| NL | 03L | Caniapiscau | 10 | 100% | 0% | 0% | Very low |
| NL | 03O | Churchill (Nfld.) | 1 | 100% | 0% | 0% | Very low |
| NL | 03M | Eastern Ungava Bay | 78 | 100% | 0% | 0% | Very low |
| NL | 02W | Gulf of St. Lawrence - Natashquan | 42 | 93% | 2% | 5% | Very low |
| NL | 02V | Gulf of St. Lawrence - Romaine | 39 | 100% | 0% | 0% | Very low |
| NL | 02U | Moisie and St. Lawrence Estuary | 61 | 100% | 0% | 0% | Very low |
| NL | 02X | Petit Mécatina and Strait of Belle Isle | 5 | 100% | 0% | 0% | Very low |
| NT | 10B | Central Liard | 14 | 43% | 36% | 21% | High |
| NT | 07L | Fond-du-Lac | 1 | 100% | 0% | 0% | Very low |
| NT | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | Very low |
| NT | 07O | Hay | 13 | 23% | 46% | 31% | Very high |
| NT | 07N | Slave | 10 | 100% | 0% | 0% | Very low |
| NT | 07P | Southern Great Slave Lake | 17 | 53% | 47% | 0% | High |
| NT | 10A | Upper Liard | 23 | 30% | 48% | 22% | High |
| ON | 04M | Abitibi | 388 | 44% | 47% | 9% | High |
| ON | 04F | Attawapiskat - Coast | 46 | 57% | 41% | 2% | High |
| ON | 02K | Central Ottawa | 1098 | 70% | 22% | 8% | High |
| ON | 02E | Eastern Georgian Bay | 752 | 91% | 4% | 5% | Very low |
| ON | 02F | Eastern Lake Huron | 186 | 5% | 12% | 82% | Very high |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-32 | >32 | Probability of Survival |
|---|---|---|---|---|---|---|---|
| ON | 05R | Eastern Lake Winnipeg | 130 | 86% | 11% | 3% | Very low |
| ON | 04E | Ekwan - Coast | 10 | 0% | 90% | 10% | High |
| ON | 05Q | English | 765 | 86% | 14% | 0% | Very low |
| ON | 20 | Great Lakes and St. Lawrence | 39 | 38% | 33% | 28% | Very high |
| ON | 04N | Harricanaw - Coast | 77 | 91% | 8% | 1% | Very low |
| ON | 04A | Hayes (Man.) | 17 | 12% | 82% | 6% | High |
| ON | 04J | Kenogami | 314 | 13% | 65% | 22% | High |
| ON | 02H | Lake Ontario and Niagara Peninsula | 539 | 55% | 27% | 18% | High |
| ON | 04H | Lower Albany - Coast | 3 | 100% | 0% | 0% | Very low |
| ON | 02L | Lower Ottawa | 503 | 82% | 14% | 4% | Very low |
| ON | 04L | Missinaibi-Mattagami | 646 | 51% | 43% | 6% | High |
| ON | 04K | Moose (Ont.) | 16 | 50% | 50% | 0% | High |
| ON | 02B | Northeastern Lake Superior | 649 | 70% | 28% | 2% | High |
| ON | 02G | Northern Lake Erie | 95 | 2% | 20% | 78% | Very high |
| ON | 02C | Northern Lake Huron | 850 | 96% | 4% | 0% | Very low |
| ON | 02A | Northwestern Lake Superior | 347 | 64% | 32% | 4% | High |
| ON | 04C | Severn | 77 | 43% | 57% | 0% | High |
| ON | 04G | Upper Albany | 229 | 72% | 27% | 1% | High |
| ON | 02J | Upper Ottawa | 854 | 89% | 10% | 1% | Very low |
| ON | 02M | Upper St. Lawrence | 89 | 25% | 51% | 25% | High |
| ON | 02D | Wanipitai and French (Ont.) | 374 | 94% | 5% | 0% | Very low |
| ON | 04D | Winisk - Coast | 45 | 42% | 56% | 2% | High |
| ON | 05P | Winnipeg | 955 | 81% | 17% | 2% | Very low |
| QC | 04M | Abitibi | 388 | 44% | 47% | 9% | High |
| QC | 02S | Betsiamites - Coast | 105 | 100% | 0% | 0% | Very low |
| QC | 03B | Broadback and Rupert | 92 | 90% | 10% | 0% | Very low |
| QC | 03L | Caniapiscau | 10 | 100% | 0% | 0% | Very low |
| QC | 02K | Central Ottawa | 1098 | 70% | 22% | 8% | High |
| QC | 02O | Central St. Lawrence | 251 | 61% | 31% | 8% | High |
| QC | 03O | Churchill (Nfld.) | 1 | 100% | 0% | 0% | Very low |
| QC | 03M | Eastern Ungava Bay | 78 | 100% | 0% | 0% | Very low |
| QC | 03C | Eastmain | 6 | 100% | 0% | 0% | Very low |
| QC | 20 | Great Lakes and St. Lawrence | 39 | 38% | 33% | 28% | Very high |
| QC | 02W | Gulf of St. Lawrence - Natashquan | 42 | 93% | 2% | 5% | Very low |
| QC | 02V | Gulf of St. Lawrence - Romaine | 39 | 100% | 0% | 0% | Very low |
| QC | 01B | Gulf of St. Lawrence and Northern Bay of Fundy (N.B.) | 37 | 19% | 59% | 22% | High |
| QC | 04N | Harricanaw - Coast | 77 | 91% | 8% | 1% | Very low |
| QC | 03D | La Grande - Coast | 1 | 100% | 0% | 0% | Very low |
| QC | 02L | Lower Ottawa | 503 | 82% | 14% | 4% | Very low |
| QC | 02P | Lower St. Lawrence | 385 | 95% | 5% | 1% | Very low |
| QC | 02T | Manicouagan and aux Outardes | 92 | 100% | 0% | 0% | Very low |
| QC | 02U | Moisie and St. Lawrence Estuary | 61 | 100% | 0% | 0% | Very low |
| QC | 02Q | Northern Gaspé Peninsula | 41 | 20% | 66% | 15% | High |
| QC | 03A | Nottaway - Coast | 209 | 97% | 3% | 0% | Very low |
| QC | 02X | Petit Mécatina and Strait of Belle Isle | 5 | 100% | 0% | 0% | Very low |
| QC | 02R | Saguenay | 343 | 99% | 1% | 0% | Very low |
| QC | 01A | Saint John and Southern Bay of | 35 | 34% | 57% | 9% | High |

| Prov. | ID | Sub-drainage | *n* | < 12 | 12-32 | >32 | Probability of Survival |
|---|---|---|---|---|---|---|---|
| | | Fundy (N.B.) | | | | | |
| QC | 02N | Saint-Maurice | 313 | 100% | 0% | 0% | Very low |
| QC | 02J | Upper Ottawa | 854 | 89% | 10% | 1% | Very low |
| QC | 02M | Upper St. Lawrence | 89 | 25% | 51% | 25% | High |
| SK | 05M | Assiniboine | 135 | 0% | 18% | 82% | Very high |
| SK | 05F | Battle | 15 | 20% | 53% | 27% | High |
| SK | 06A | Beaver (Alta.-Sask.) | 47 | 6% | 74% | 19% | High |
| SK | 07C | Central Athabasca - Lower | 19 | 21% | 63% | 16% | High |
| SK | 06E | Central Churchill (Man.) - Lower | 68 | 50% | 19% | 31% | Very high |
| SK | 06C | Central Churchill (Man.) - Upper | 1 | 0% | 100% | 0% | High |
| SK | 05E | Central North Saskatchewan | 45 | 11% | 62% | 27% | Very high |
| SK | 07L | Fond-du-Lac | 1 | 100% | 0% | 0% | Very low |
| SK | 07Q | Great Slave Lake - East Arm South Shore | 2 | 100% | 0% | 0% | Very low |
| SK | 71 | Lake Athabasca | 2 | 50% | 50% | 0% | High |
| SK | 07M | Lake Athabasca - Shores | 6 | 100% | 0% | 0% | Very low |
| SK | 05L | Lake Winnipegosis and Lake Manitoba | 105 | 0% | 7% | 93% | Very high |
| SK | 07D | Lower Athabasca | 37 | 41% | 51% | 8% | High |
| SK | 05G | Lower North Saskatchewan | 5 | 40% | 40% | 20% | High |
| SK | 05H | Lower South Saskatchewan | 42 | 0% | 0% | 100% | Very high |
| SK | 11A | Missouri | 3 | 0% | 67% | 33% | Very high |
| SK | 05J | Qu'Appelle | 40 | 0% | 3% | 98% | Very high |
| SK | 05C | Red Deer | 35 | 14% | 37% | 49% | Very high |
| SK | 06D | Reindeer | 1 | 100% | 0% | 0% | Very low |
| SK | 05K | Saskatchewan | 156 | 0% | 48% | 52% | Very high |
| SK | 05N | Souris | 35 | 0% | 9% | 91% | Very high |
| SK | 06B | Upper Churchill (Man.) | 1 | 0% | 100% | 0% | High |
| SK | 05A | Upper South Saskatchewan | 52 | 0% | 42% | 58% | Very high |
| YT | 08A | Alsek | 5 | 0% | 60% | 40% | Very high |
| YT | 10B | Central Liard | 14 | 43% | 36% | 21% | High |
| YT | 09A | Headwaters Yukon | 38 | 61% | 37% | 3% | High |
| YT | 10A | Upper Liard | 23 | 30% | 48% | 22% | High |

*Table 13. Probability of Quagga Mussel arrival, survival, and invasion per sub-drainage. The probability of invasion is based on the probability of survival (calcium suitability) and probability of arrival (propagule pressure corrected for proximity to an invaded watershed). The risk to the environment is based on the probability of invasion and impacts to the environment.*

| Prov. | ID | Sub-drainage | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|
| AB | 05F | Battle | High | High | 0 | High | High | High |
| AB | 06A | Beaver (Alta.-Sask.) | High | Moderate | 0 | Moderate | High | High |
| AB | 05B | Bow | Very high | High | 0 | High | Very high | High |
| AB | 07C | Central Athabasca - Lower | High | Low | 0 | Low | Moderate | High |
| AB | 07B | Central Athabasca - Upper | High | Moderate | 0 | Moderate | High | High |
| AB | 05E | Central North Saskatchewan | Very high | High | 0 | High | Very high | High |
| AB | 07J | Central Peace - Lower | High | Low | 0 | Low | Moderate | High |
| AB | 07H | Central Peace - Upper | Very high | Low | 0 | Low | High | High |
| AB | 08N | Columbia - U.S.A. | Very high | Moderate | 0 | Moderate | High | High |
| AB | 10C | Fort Nelson | Very high | Low | 0 | Low | High | High |
| AB | 07Q | Great Slave Lake - East Arm South Shore | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 07O | Hay | Very high | Low | 0 | Low | High | High |
| AB | 71 | Lake Athabasca | High | Very low | 0 | Very low | Moderate | High |
| AB | 07M | Lake Athabasca - Shores | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 07D | Lower Athabasca | High | Very low | 0 | Very low | Moderate | High |
| AB | 05G | Lower North Saskatchewan | High | High | 0 | High | High | High |
| AB | 07K | Lower Peace | Very low | Very low | 0 | Very low | Very low | Low |
| AB | 05H | Lower South Saskatchewan | Very high | High | 0 | High | Very high | High |
| AB | 11A | Missouri | Very high | Moderate | 0 | Moderate | High | High |
| AB | 05C | Red Deer | Very high | High | 0 | High | Very high | High |
| AB | 07N | Slave | Very low | Low | 0 | Low | Very low | Low |
| AB | 07G | Smoky | Very high | Moderate | 0 | Moderate | High | High |
| AB | 07P | Southern Great Slave Lake | High | Very low | 0 | Very low | Moderate | High |
| AB | 07A | Upper Athabasca | Very high | Moderate | 0 | Moderate | High | High |
| AB | 06B | Upper Churchill (Man.) | High | Very low | 0 | Very low | Moderate | High |
| AB | 08K | Upper Fraser | Very high | Low | 0 | Low | High | High |
| AB | 05D | Upper North Saskatchewan | High | Moderate | 0 | Moderate | High | High |
| AB | 07F | Upper Peace | Very high | Moderate | 0 | Moderate | High | High |
| AB | 05A | Upper South Saskatchewan | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08A | Alsek | Very high | Low | 0 | Low | High | High |
| BC | 05B | Bow | Very high | High | 0 | High | Very high | High |

| Prov. | ID | Sub-drainage | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|
| BC | 08F | Central Coastal Waters of B.C. | Very low | Low | 0 | Low | Very low | Low |
| BC | 10B | Central Liard | High | Very low | 0 | Very low | Moderate | High |
| BC | 08N | Columbia - U.S.A. | Very high | Moderate | 0 | Moderate | High | High |
| BC | 10C | Fort Nelson | Very high | Low | 0 | Low | High | High |
| BC | 07O | Hay | Very high | Low | 0 | Low | High | High |
| BC | 09A | Headwaters Yukon | High | Low | 0 | Low | Moderate | High |
| BC | 08M | Lower Fraser | High | Moderate | 0 | Moderate | High | High |
| BC | 08D | Nass - Coast | High | Low | 0 | Low | Moderate | High |
| BC | 08J | Nechako | High | Low | 0 | Low | Moderate | High |
| BC | 08B | Northern Coastal Waters of B.C. | High | Low | 0 | Low | Moderate | High |
| BC | 08O | Queen Charlotte Islands | Very low | Moderate | 0 | Moderate | Very low | Low |
| BC | 08E | Skeena - Coast | High | Low | 0 | Low | Moderate | High |
| BC | 07G | Smoky | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08G | Southern Coastal Waters of B.C. | Very low | Low | 0 | Low | Very low | Low |
| BC | 08C | Stikine - Coast | High | Low | 0 | Low | Moderate | High |
| BC | 08L | Thompson | Very high | Moderate | 0 | Moderate | High | High |
| BC | 07A | Upper Athabasca | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08K | Upper Fraser | Very high | Low | 0 | Low | High | High |
| BC | 10A | Upper Liard | High | Low | 0 | Low | Moderate | High |
| BC | 05D | Upper North Saskatchewan | High | Moderate | 0 | Moderate | High | High |
| BC | 07F | Upper Peace | Very high | Moderate | 0 | Moderate | High | High |
| BC | 05A | Upper South Saskatchewan | Very high | Moderate | 0 | Moderate | High | High |
| BC | 08H | Vancouver Island | Very low | Moderate | 0 | Moderate | Very low | Low |
| BC | 07E | Williston Lake | High | Low | 0 | Low | Moderate | High |
| MB | 05M | Assiniboine | Very high | High | 0 | High | Very high | High |
| MB | 06E | Central Churchill (Man.) - Lower | Very high | Very low | 0 | Very low | Moderate | High |
| MB | 05R | Eastern Lake Winnipeg | Very low | Very low | 1 | Low | Very low | Low |
| MB | 05T | Grass and Burntwood | High | Low | 0 | Low | Moderate | High |
| MB | 04A | Hayes (Man.) | High | Very low | 0 | Very low | Moderate | High |
| MB | 50 | Lake Winnipeg | Very high | Low | 0 | Low | High | High |
| MB | 05L | Lake Winnipegosis and Lake Manitoba | Very high | Moderate | 0 | Moderate | High | High |
| MB | 06F | Lower Churchill (Man.) | High | Very low | 0 | Very low | Moderate | High |
| MB | 05U | Nelson | Very high | Very low | 0 | Very low | Moderate | High |
| MB | 05J | Qu'Appelle | Very high | High | 0 | High | Very high | High |
| MB | 05O | Red | Very high | High | 1 | Very high | Very high | High |

| Prov. | ID | Sub-drainage | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|
| MB | 06D | Reindeer | Very low | Very low | 0 | Very low | Very low | Low |
| MB | 05K | Saskatchewan | Very high | Moderate | 0 | Moderate | High | High |
| MB | 04C | Severn | High | Very low | 0 | Very low | Moderate | High |
| MB | 05N | Souris | Very high | High | 0 | High | Very high | High |
| MB | 05S | Western Lake Winnipeg | Very high | Moderate | 1 | High | Very high | High |
| MB | 05P | Winnipeg | Very low | Moderate | 1 | High | Very low | Low |
| NL | 03L | Caniapiscau | Very low | Very low | 1 | Low | Very low | Low |
| NL | 03O | Churchill (Nfld.) | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 03M | Eastern Ungava Bay | Very low | Low | 0 | Low | Very low | Low |
| NL | 02W | Gulf of St. Lawrence - Natashquan | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 02V | Gulf of St. Lawrence - Romaine | Very low | Very low | 1 | Low | Very low | Low |
| NL | 02U | Moisie and St. Lawrence Estuary | Very low | Low | 1 | Moderate | Very low | Low |
| NL | 02X | Petit Mécatina and Strait of Belle Isle | Very low | Low | 0 | Low | Very low | Low |
| NT | 10B | Central Liard | High | Very low | 0 | Very low | Moderate | High |
| NT | 07L | Fond-du-Lac | Very low | Very low | 0 | Very low | Very low | Low |
| NT | 07Q | Great Slave Lake - East Arm South Shore | Very low | Very low | 0 | Very low | Very low | Low |
| NT | 07O | Hay | Very high | Low | 0 | Low | High | High |
| NT | 07N | Slave | Very low | Low | 0 | Low | Very low | Low |
| NT | 07P | Southern Great Slave Lake | High | Very low | 0 | Very low | Moderate | High |
| NT | 10A | Upper Liard | High | Low | 0 | Low | Moderate | High |
| ON | 04M | Abitibi | High | Moderate | 1 | High | High | High |
| ON | 04F | Attawapiskat - Coast | High | Very low | 0 | Very low | Moderate | High |
| ON | 02K | Central Ottawa | High | Moderate | 1 | High | High | High |
| ON | 02E | Eastern Georgian Bay | Very low | High | 1 | Very high | Very low | Low |
| ON | 02F | Eastern Lake Huron | Very high | Very high | 1 | Very high | Very high | High |
| ON | 05R | Eastern Lake Winnipeg | Very low | Very low | 1 | Low | Very low | Low |
| ON | 04E | Ekwan - Coast | High | Very low | 0 | Very low | Moderate | High |
| ON | 05Q | English | Very low | Low | 1 | Moderate | Very low | Low |
| ON | 20 | Great Lakes and St. Lawrence | Very high | Very high | 1 | Very high | Very high | High |
| ON | 04N | Harricanaw - Coast | Very low | Low | 1 | Moderate | Very low | Low |
| ON | 04A | Hayes (Man.) | High | Very low | 0 | Very low | Moderate | High |
| ON | 04J | Kenogami | High | Low | 1 | Moderate | High | High |
| ON | 02H | Lake Ontario and Niagara Peninsula | High | Very high | 1 | Very high | Very high | High |
| ON | 04H | Lower Albany - Coast | Very low | Very low | 0 | Very low | Very low | Low |
| ON | 02L | Lower Ottawa | Very low | High | 1 | Very high | Very low | Low |

| Prov. | ID | Sub-drainage | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|
| ON | 04L | Missinaibi-Mattagami | High | Moderate | 1 | High | High | High |
| ON | 04K | Moose (Ont.) | High | Very low | 1 | Low | Moderate | High |
| ON | 02B | Northeastern Lake Superior | High | Low | 1 | Moderate | High | High |
| ON | 02G | Northern Lake Erie | Very high | Very high | 1 | Very high | Very high | High |
| ON | 02C | Northern Lake Huron | Very low | Moderate | 1 | High | Very low | Low |
| ON | 02A | Northwestern Lake Superior | High | Moderate | 1 | High | High | High |
| ON | 04C | Severn | High | Very low | 0 | Very low | Moderate | High |
| ON | 04G | Upper Albany | High | Very low | 1 | Low | Moderate | High |
| ON | 02J | Upper Ottawa | Very low | Moderate | 1 | High | Very low | Low |
| ON | 02M | Upper St. Lawrence | High | High | 1 | Very high | Very high | High |
| ON | 02D | Wanipitai and French (Ont.) | Very low | Moderate | 1 | High | Very low | Low |
| ON | 04D | Winisk - Coast | High | Very low | 0 | Very low | Moderate | High |
| ON | 05P | Winnipeg | Very low | Moderate | 1 | High | Very low | Low |
| QC | 04M | Abitibi | High | Moderate | 1 | High | High | High |
| QC | 02S | Betsiamites - Coast | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03B | Broadback and Rupert | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03L | Caniapiscau | Very low | Very low | 1 | Low | Very low | Low |
| QC | 02K | Central Ottawa | High | Moderate | 1 | High | High | High |
| QC | 02O | Central St. Lawrence | High | Very high | 1 | Very high | Very high | High |
| QC | 03O | Churchill (Nfld.) | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03M | Eastern Ungava Bay | Very low | Low | 0 | Low | Very low | Low |
| QC | 03C | Eastmain | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 20 | Great Lakes and St. Lawrence | Very high | Very high | 1 | Very high | Very high | High |
| QC | 02W | Gulf of St. Lawrence - Natashquan | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02V | Gulf of St. Lawrence - Romaine | Very low | Very low | 1 | Low | Very low | Low |
| QC | 01B | Gulf of St. Lawrence and Northern Bay of Fundy (N.B.) | High | High | 1 | Very high | Very high | High |
| QC | 04N | Harricanaw - Coast | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 03D | La Grande - Coast | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02L | Lower Ottawa | Very low | High | 1 | Very high | Very low | Low |
| QC | 02P | Lower St. Lawrence | Very low | High | 1 | Very high | Very low | Low |
| QC | 02T | Manicouagan and aux Outardes | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02U | Moisie and St. Lawrence Estuary | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02Q | Northern Gaspé Peninsula | High | High | 1 | Very high | Very high | High |
| QC | 03A | Nottaway - Coast | Very low | Low | 1 | Moderate | Very low | Low |
| QC | 02X | Petit Mécatina and Strait of Belle Isle | Very low | Low | 0 | Low | Very low | Low |

| Prov. | ID | Sub-drainage | Probability of Survival | Propagule Pressure | Prox corr. | Probability of Arrival | Probability of Invasion | Risk to Environment |
|---|---|---|---|---|---|---|---|---|
| QC | 02R | Saguenay | Very low | Moderate | 1 | High | Very low | Low |
| QC | 01A | Saint John and Southern Bay of Fundy (N.B.) | High | High | 1 | Very high | Very high | High |
| QC | 02N | Saint-Maurice | Very low | Moderate | 1 | High | Very low | Low |
| QC | 02J | Upper Ottawa | Very low | Moderate | 1 | High | Very low | Low |
| QC | 02M | Upper St. Lawrence | High | High | 1 | Very high | Very high | High |
| SK | 05M | Assiniboine | Very high | High | 0 | High | Very high | High |
| SK | 05F | Battle | High | High | 0 | High | High | High |
| SK | 06A | Beaver (Alta.-Sask.) | High | Moderate | 0 | Moderate | High | High |
| SK | 07C | Central Athabasca - Lower | High | Low | 0 | Low | Moderate | High |
| SK | 06E | Central Churchill (Man.) - Lower | Very high | Very low | 0 | Very low | Moderate | High |
| SK | 06C | Central Churchill (Man.) - Upper | High | Low | 0 | Low | Moderate | High |
| SK | 05E | Central North Saskatchewan | Very high | High | 0 | High | Very high | High |
| SK | 07L | Fond-du-Lac | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 07Q | Great Slave Lake - East Arm South Shore | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 71 | Lake Athabasca | High | Very low | 0 | Very low | Moderate | High |
| SK | 07M | Lake Athabasca - Shores | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 05L | Lake Winnipegosis and Lake Manitoba | Very high | Moderate | 0 | Moderate | High | High |
| SK | 07D | Lower Athabasca | High | Very low | 0 | Very low | Moderate | High |
| SK | 05G | Lower North Saskatchewan | High | High | 0 | High | High | High |
| SK | 05H | Lower South Saskatchewan | Very high | High | 0 | High | Very high | High |
| SK | 11A | Missouri | Very high | Moderate | 0 | Moderate | High | High |
| SK | 05J | Qu'Appelle | Very high | High | 0 | High | Very high | High |
| SK | 05C | Red Deer | Very high | High | 0 | High | Very high | High |
| SK | 06D | Reindeer | Very low | Very low | 0 | Very low | Very low | Low |
| SK | 05K | Saskatchewan | Very high | Moderate | 0 | Moderate | High | High |
| SK | 05N | Souris | Very high | High | 0 | High | Very high | High |
| SK | 06B | Upper Churchill (Man.) | High | Very low | 0 | Very low | Moderate | High |
| SK | 05A | Upper South Saskatchewan | Very high | Moderate | 0 | Moderate | High | High |
| YT | 08A | Alsek | Very high | Low | 0 | Low | High | High |
| YT | 10B | Central Liard | High | Very low | 0 | Very low | Moderate | High |
| YT | 09A | Headwaters Yukon | High | Low | 0 | Low | Moderate | High |
| YT | 10A | Upper Liard | High | Low | 0 | Low | Moderate | High |

## FIGURES




*Figure 1. Freshwater drainages used in the risk assessment of three dreissenid mussels. Sub-drainages are identified by three-digit identifiers and hatched watersheds were excluded. See Appendices A4 and A5 for close-ups of provinces. Source: Natural Resources Canada.*




*Figure 2. Quagga Mussels (Dreissena rostriformis bugensis) (1-6) and Zebra Mussels (Dreissena polymorpha) (7-9) collected from the River Main (Germany). From: Van der Velde, G. and Platvoet, D. 2007*




*Figure 3. Comparison of shell structures of the Zebra Mussel (Dreissena polymporha) and the Quagga Mussel (D. rostriformis bugensis). Source: United States Geological Survey*

*Figure 4. Comparison of Zebra Mussel (upper) and Dark Falsemussel (lower) shells. From: Verween, A., Vincx, M. and Degraer, S. 2010.*




*Figure 5. Framework for the restructuring of food webs by Zebra Mussel and Quagga Mussel. Arrows represent the direction of energy flow. Bold lines and plus symbols (+) represent increased fluxes, while minus symbols (-) represent decreased fluxes. Modified from Higgins (in press).*




*Figure 6. Reported sightings of adult Zebra Mussels between 1986 and 2011 in North America. Data obtained from the US Geological Society (Benson et al., 2012a).*




*Figure 7. Reported sightings of adult Quagga Mussels between 1986 and 2011 in North America. Data obtained from the US Geological Society (Benson et al., 2012b).*




*Figure 8. Flow diagram of risk assessment process for Zebra Mussel and Quagga Mussel invasion into Canadian Freshwaters.*

| Impact to Environment | | | | | | |
|---|---|---|---|---|---|---|
| | Very High | | | | | |
| | High | | | | | |
| | Moderate | | | | | |
| | Low | | | | | |
| | Very Low | | | | | |
| | | Very Low | Low | Moderate | High | Very High |
| | Probability of Invasion | | | | | |

*Figure 9. Heat (risk) matrix used to determine the ecological risk posed by three dreissenid species for six Canadian provinces (BC, AB, SK, MB, ON, QC). The risk to the environment is calculated by combining the probability of invasion determined in Step 1 with the impacts of invasion determined in Step 2. In the matrix risk to the environment is: Green = Low Risk, Yellow = Moderate Risk, and Red = High Risk.*




*Figure 10. Calcium concentrations (mg/L, 75th percentile) per sub-drainage. Sub-drainageware ranked on their suitability for Zebra Mussel survival based on calcium concentrations required to develop their shells: (< 12 mg/L = below (pale green); 12-19 mg/L = moderate (yellow); 20-25 mg/L = high (orange); > 25 mg/L = very high (red)). See Annex A1 for calcium concentions per sub-sub-drainage. Hatched watersheds had less than 5 sampling sites.*

*Figure 11. Temperature correction factor based on mean air temperature of the warmest quarter per sub-drainage where red colored sub-drainages had calcium suitability reduced by one category. Data obtained from WorldClim (http://www.worldclim.org/)*

*Figure 12. Probability of survival (habitat suitability) for Zebra Mussel per sub-drainage based on calcium concentrations (mg/L, 75th percentile) and corrected for temperature. Hatched watersheds had less than 5 sampling sites.*

*Figure 13. Propagule pressure derived from the Human Footprint Index per Canadian sub-drainage (modified from Sanderson et al. 2002). The index is a composite factor of human influence corrected by biome type that integrates data of land use, urbanization, population density, transportation networks and other human activities that are known to facilitate species invasions. Indices are color coded from low (light green) to very high (red). See Annex #3 for original data.*




*Figure 14. Proximity to invaded Zebra Mussel sub-drainages.*




*Figure 15. Probability of Zebra Mussel arrival per assessed sub-drainage based on propagule pressure and proximity to invaded sub-drainages. The probability of arrival is scored as very low (pale green) to very high (red).*




*Figure 16. Probability of Zebra Mussel invasion based on probability of survival and arrival. Hatched watersheds had less than 5 sampling sites.*




*Figure 17. Zebra Mussel ecological risk per sub-drainage, ranging from low to high. Risk is based on probability of invasion and impacts on the environment. Hatched watersheds had less than 5 sampling sites.*




*Figure 18. Probability of survival (habitat suitability) for Quagga Mussel per sub-drainage based on calcium concentrations (mg/L, 75th percentile). Sub-drainages were ranked on their suitability for Quagga Mussel survival based on literature accounts (see Table 1). Results per sub-sub-drainage can be found in Annex A2. Hatched watersheds had less than 5 sampling sites.*




*Figure 19. Affected and adjoining Canadian sub-drainages where Quagga Mussel sightings have been reported in North America.*




*Figure 20. Probability of Quagga Mussel arrival per assessed sub-drainage based on propagule pressure and proximity to invaded sub-drainages. The probability of arrival is scored as very low (pale green) to very high (red).*




*Figure 21. Probability of Quagga Mussel invasion based on probability of survival and arrival. Hatched watersheds had less than 5 sampling sites.*




*Figure 22. Quagga Mussel ecological risk per sub-drainage, ranging from low to high. Risk is based on probability of invasion and impacts on the environment. Hatched watersheds had less than 5 sampling sites.*




*Figure A1. Calcium concentrations (mg/L, 75th percentile) per sub-sub-drainage. Sub-sub-drainages were ranked on their suitability for Zebra Mussel survival based on literature accounts: (< 12 mg/L = very low (pale green); 12-19 mg/L = moderate (orange); 20-25 mg/L = high (orange); > 25 mg/L = very high (red)). Hatched sub-drainages had less than 5 sampling sites.*




*Figure A2. Calcium concentrations (mg/L, 75th percentile) per sub-sub-drainage. Sub-sub-drainages were ranked on their suitability for Quagga Mussel survival based on literature accounts: (< 12 mg/L = very low (pale green); 12-32 mg/L = high (orange); > 32 mg/L = very high (red)). Hatched watersheds had less than 5 sampling sites.*




*Figure A3. The Human Footprint Index (Sanderson et al. 2002; see http://sedac.ciesin.columbia.edu/wildareas/), is a composite factor of human influence corrected by biome type that integrates data of land use, urbanization, population density, transportation networks and other human activities that are known to facilitate species invasions.*




*Figure A4. Freshwater sub-drainages in British-Columbia, Alberta, and Saskatchewan. Sub-drainages are identified by three-digit identifiers. Source: Natural Resources Canada.*




*Figure A5. Freshwater sub-drainages in Manitoba, Ontario, and Quebec. Sub-drainages are identified by three-digit identifiers. Source: Natural Resources Canada.*